

2015

Independent evidence for an association between general cognitive ability and a genetic locus for educational attainment

J. W. Trampush

Hofstra Northwell School of Medicine

T. Lencz

Hofstra Northwell School of Medicine

S. Guha

Northwell Health

S. Mukherjee

Northwell Health

P. DeRosse

Northwell Health

See next page for additional authors

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/articles>



Part of the [Psychiatry Commons](#)

Recommended Citation

Trampush JW, Lencz T, Guha S, Mukherjee S, DeRosse P, John M, Andreassen O, Deary I, Glahn D, Malhotra AK, . Independent evidence for an association between general cognitive ability and a genetic locus for educational attainment. . 2015 Jan 01; 168(5):Article 927 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/articles/927>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works.

Authors

J. W. Trampush, T. Lencz, S. Guha, S. Mukherjee, P. DeRosse, M. John, O. A. Andreassen, I. J. Deary, D. C. Glahn, A. K. Malhotra, and +41 additional authors



HHS Public Access

Author manuscript

Am J Med Genet B Neuropsychiatr Genet. Author manuscript; available in PMC 2016 July 01.

Published in final edited form as:

Am J Med Genet B Neuropsychiatr Genet. 2015 July ; 168(5): 363–373. doi:10.1002/ajmg.b.32319.

Independent Evidence for an Association between General Cognitive Ability and a Genetic Locus for Educational Attainment

A full list of authors and affiliations appears at the end of the article.

Abstract

Cognitive deficits and reduced educational achievement are common in psychiatric illness; understanding the genetic basis of cognitive and educational deficits may be informative about the etiology of psychiatric disorders. A recent, large genomewide association study (GWAS) reported a genome-wide significant locus for years of education, which subsequently demonstrated association to general cognitive ability (“g”) in overlapping cohorts. The current study was designed to test whether GWAS hits for educational attainment are involved in general cognitive ability in an independent, large-scale collection of cohorts. Using cohorts in the Cognitive Genomics Consortium (COGENT; up to 20,495 healthy individuals), we examined the relationship between g and variants associated with educational attainment. We next conducted meta-analyses with 24,189 individuals with neurocognitive data from the educational attainment studies, and then with 53,188 largely independent individuals from a recent GWAS of cognition. A SNP (rs1906252) located at chromosome 6q16.1, previously associated with years of schooling, was significantly associated with g ($P = 1.47 \times 10^{-4}$) in COGENT. The first joint analysis of 43,381 non-overlapping individuals for this a priori-designated locus was strongly significant ($P = 4.94 \times 10^{-7}$), and the second joint analysis of 68,159 non-overlapping individuals was even more robust ($P = 1.65 \times 10^{-9}$). These results provide independent replication, in a large-scale dataset, of a genetic locus associated with cognitive function and education. As sample sizes grow, cognitive GWAS will identify increasing numbers of associated loci, as has been accomplished in other polygenic quantitative traits, which may be relevant to psychiatric illness.

Keywords

neurocognition; general cognitive ability; educational attainment; genetics; GWAS; proxy phenotype

*Correspondence: Joey W. Trampush, Zucker Hillside Hospital, Division of Psychiatry Research, 75-59 263rd Street, Glen Oaks, NY, 11004, USA, jtrampush@nshs.edu.

Description of Supplemental Data

Supplemental Data include detailed descriptions of the 21 COGENT cohorts including neurocognitive and genetic assays employed across sites. In addition, detailed statistical methodology, imputation methods, and quality control procedures of phenotypes and genotypes are described in detail, and three supplemental figures along with two supplemental tables and supplemental references are included.

Competing Financial Interests

The authors declare no conflict of interest.

Introduction

A general cognitive ability factor (also termed *g*) typically captures just under half of the overall variance in performance on diverse laboratory measures of neurocognitive functioning. (Johnson et al., 2008) General performance on neurocognitive tests has remarkable predictive value across a diverse range of social, health and behavioral outcomes, more so than any other psychological trait. (Gottfredson, 1997; Deary et al., 2011; Deary, 2012) As examples, low *g* performance is associated with lower educational attainment and income, (Johnson et al., 2009) is a better predictor of mortality from cardiovascular disease than smoking, blood glucose and cholesterol, (Deary, 2008) and predicts longevity. (Batty et al., 2008) Deficits in general neurocognitive performance are pervasive in most psychiatric and neurologic disorders, yet are often the most difficult component to treat. (Millan et al., 2012) As such, understanding the neurobiology of human cognition is potentially critical to improving physical and mental health outcomes in society. (Deary et al., 2010)

While both genetic background and environmental experience interact to shape cognitive development, (Deary et al., 2012) twin and family studies have consistently demonstrated heritability of more than 50% for general cognitive ability measured in adulthood. (Deary et al., 2009) Allelic variation can have a direct influence on brain biology by modifying the molecular structure and/or function of brain-expressed transcripts and proteins such as neurotransmitter receptors and neurodevelopmental growth factors. (Chen et al., 2004) However, attempts to pinpoint loci associated with human cognition across diverse population samples have proven challenging due to the difficulty of assembling the large cohorts required to detect small expected effects of individual variants in a highly polygenic trait. (Benyamin et al., 2014; Chabris et al., 2012; Luciano et al., 2011; Lencz et al., 2013; Need et al., 2009; Davies et al., 2011, 2015; Martin et al., 2011)

By contrast, *educational history* is easily obtainable demographic information collected in any field of medical research, and can therefore be collected in more readily across large cohorts as compared to cognition. Educational attainment, as measured by self-reported years of schooling, has been proposed as a 'proxy phenotype' for cognitive ability for GWAS since much larger samples can be utilized compared to neurocognitive studies. (Rietveld et al., 2014a, 2014b, 2013; Martin et al., 2011) The Social Science Genetic Association Consortium (SSGAC) reported on a 126,559 person GWAS that detected three genome-wide significant SNPs associated with completion of college (rs11584700 and rs4851266) and years of schooling (rs9320913). (Rietveld et al., 2013) In a post hoc analysis, these SNPs had a stronger and more direct effect on cognitive function than on education. (Rietveld et al., 2013) Further, a polygenic risk score of educational attainment SNPs accounted for 2-3% of the variance in general cognitive ability in an independent sample, and a mediation analysis suggested that *g* mediated more than half of the effect these SNPs had on education. (Rietveld et al., 2013) Here, we analyzed the three SNPs obtained in the SSGAC educational attainment GWAS in $\approx 20,000$ independent subjects in the Cognitive Genomics Consortium (COGENT), (Donohoe et al., 2012; Lencz et al., 2013) and found converging evidence across multiple large cohorts that common variation at genomic region 6q16.1, previously associated with years of schooling, reliably predicts variation in *g*.

Methods and Materials

COGENT is an international GWAS collaboration formed to conduct genetic analyses of g and related neurocognitive processes in healthy individuals.(Donohoe et al., 2012) Though common GWAS markers have been proposed to account for ~30% or more of the variance in general intelligence in adults, individual SNPs only contribute a small fraction of the variance to the heritability of g due to extreme polygenicity.(Davies et al., 2011; Marioni et al., 2014) Detecting SNP associations of such small magnitudes via GWAS requires large samples many times the size an individual lab can ascertain, leading to consortia such as COGENT. The decision to study g in COGENT stemmed from longstanding evidence that a g factor can be derived consistently, captures almost half the variance in overall test performance, and is relatively invariant to the neurocognitive test battery used and specific abilities assessed.(Johnson et al., 2008; Panizzon et al., 2014)

The first phase of COGENT ('COGENT1') resulted in a GWAS of general cognitive ability in $\approx 5,000$ individuals from the general population.(Lencz et al., 2013) The next (and ongoing) wave of data collection in COGENT ('COGENT2') has resulted in the acquisition of $\approx 15,000$ independent subjects with neurocognitive and GWAS data for analysis (see Table 1 for cohort details). To be included as a participant in COGENT, data from at least one neuropsychological measure across at least three domains of cognitive performance (e.g., digit span for working memory; logical memory for verbal declarative memory; and digit symbol coding for processing speed), or the use of a validated g -sensitive measure was required. Tests missing for more than 5% of the sample in an individual study were excluded. Each COGENT study administered an average of 10.3 neurocognitive tests, and the internal consistency of performance within each study was strong (mean Cronbach's $\alpha = 75\%$; supplementary table S1). The first unrotated principal component accounted for just under half of the variance across the 21 studies on average, as expected based on an extensive prior literature.(Carroll, 1993) As Figure 1 shows, g had normal distributional properties within all 21 studies, a feature critical to enhancing statistical power in quantitative trait analysis.

All COGENT samples were genotyped on commercial Illumina or Affymetrix genome-wide SNP microarrays (supplementary table S1). Standard GWAS quality control (QC) methods were applied to the genetic data (described in detail in the supplementary information). Subjects in the study were Caucasian of European ancestry, which we confirmed by analysis of genotype data using multidimensional scaling (MDS). Genetic outliers were removed in each study based on MDS axis plotting versus HapMap3 ethnic subgroups. Note that none of the three SNPs previously associated with education were variants included on commercially available microarrays, and thus were imputed into their datasets.(Rietveld et al., 2013) In the COGENT1 studies, SNPs were imputed using HapMap3 reference panels as previously described.(Lencz et al., 2013) COGENT2 samples that did not have genotypes for the SNPs of interest were imputed using IMPUTE2(Howie et al., 2009) and 1000 Genomes Project reference panels (downloaded June 2014).

SNP analysis of g was completed separately within each COGENT study using Plink(Purcell et al., 2007) or Genome-wide Complex Trait Analysis (GCTA).(Yang et al.,

2011) Plink was used to analyze datasets comprising unrelated individuals, and GCTA used to analyze five datasets in which multiple family members were known to be included *a priori*. GCTA has implemented a mixed-linear-model association (MLMA) analytic function that corrects for population or relatedness structure through a correction that is specific to the structure of interest.(Yang et al., 2014) Regression coefficients and standard error estimates were generated within each study and then carried forward for meta-analysis using Metasoft(Han and Eskin, 2011) and the R MetABEL package(Aulchenko et al., 2007) for plotting of results. Although we expected that allelic effects for cognitive ability would mirror the direction observed for educational attainment, all analyses were conservatively carried out using two-tailed tests.

A multi-stage approach was utilized to determine if GWAS hits associated with educational attainment were also associated with general cognitive ability in COGENT. First, we examined the *p*-values of the three educational attainment SNPs (and their close proxies) in the database housing the results of the COGENT1 GWAS.(Lencz et al., 2013) SNPs with *p*-values < .05 identified in this first stage were then meta-analyzed for association to *g* in ≈16,000 independent subjects in the subsequent replication stage.

Results

COGENT Analysis

Using the approach described above, the two SNPs previously associated with the dichotomous variable of completing college (rs11584700 and rs4851266) in(Rietveld et al., 2013) were not significantly associated with *g* in COGENT1 (*P*'s > .05). The third SNP (rs9320913) associated with years of schooling was neither genotyped nor imputed in COGENT1; however, a close proxy SNP (rs1906252, $R^2 = 1.0$ in HapMap2 CEU, $R^2 = .905$ in 1000 Genomes CEU) was available for analysis in the COGENT1 GWAS data. rs1906252 was either typed or imputed in all nine COGENT1 studies, and was significantly associated with general cognitive ability in the expected direction, such that the minor (A) allele was associated with higher *g* scores ($\beta = .050$, $P = 1.20 \times 10^{-2}$; $P_Q = .340$). This SNP was therefore carried forward for analysis in COGENT2.

To date, GWAS and neurocognitive data have been acquired for more than 15,000 participants from 12 independent cohorts in COGENT2. The average sample size of individual COGENT2 studies was 1,398 and ranged from 127 to 4,412 Caucasian subjects of European ancestry. rs1906252 was genotyped or imputed in all 12 COGENT2 samples (N = 16,544 genotypes), of which 15,533 had *g* factor data. As shown in Table 2, the association between allelic variation at SNP rs1906252 and *g* in COGENT2 was statistically significant and not biased by heterogeneity ($\beta = .027$, $P = 2.34 \times 10^{-3}$, two-tailed; $P_Q = .671$). The direction of the minor allele effect was positive in 10 out of 12 COGENT2 studies, which was consistent with the educational attainment GWAS studies and COGENT1, and statistically different from chance (binomial test, $P = .02$). Finally, all 21 COGENT datasets were merged to analyze the association between rs1906252 and *g* in the combined sample of 20,495 individuals. A significant association was again detected in the combined analysis that was not confounded by heterogeneity ($\beta = .031$, $P = 1.47 \times 10^{-4}$, two-tailed; $P_Q = .543$). Figure 2 presents the combined results in a forest plot, and as shown

in Figure 3, a linear increase in general cognitive ability was related positively to minor A allele load.

We tested the robustness of the association between rs1906252 and *g* in a series of sensitivity analyses. First, a potential confound worthy of consideration was the fact that two COGENT1 studies (Helsinki Birth Cohort Study [HBCS; $n = 332$] and the 1936 Lothian Birth Cohort Study [LBC1936; $n = 1,005$]) (Lencz et al., 2013) were also included in SSGAC. (Rietveld et al., 2014b, 2013) The non-independence of our collective studies and the high correlation between educational attainment and cognition necessitated a sensitivity analysis requiring HBCS and LBC1936 be excluded and the remaining COGENT studies reanalyzed. This analysis also yielded a significant result between variation in rs1906252 and *g* ($\beta = .026$, $P = 1.60 \times 10^{-3}$, two-tailed; $P_Q = .889$; see Table 2).

Next, age had a bimodal distribution across the 21 COGENT studies. The first peak was at approximately 18 years of age, and the second peak at approximately 70 years (supplemental figure S1). To explore the effect of age on the association between rs1906252 and *g* across the lifespan, we examined the interaction between rs1906252 and age on *g*, which was not significant ($\beta = .0004$, $P = .302$). We then split the sample at the ‘valley’ of the distribution at age 40. This resulted in a group of 7,208 individuals under 40 years of age and a group of 13,287 individuals 40 years of age or older. The interaction between rs1906252 and this dichotomous age split was not significant ($\beta = .007$, $P = .728$). Thus, age did not moderate the association between rs1906252 and general cognitive function across the lifespan in COGENT.

SSGAC Meta-Analysis

The SSGAC reanalyzed their GWAS data utilizing a two-stage design in order to examine whether educational attainment is a valid proxy phenotype for cognitive ability. (Rietveld et al., 2014b) Subjects with available neurocognitive data ($n = 24,189$) were removed from the larger cohort, and a GWAS of years of education was then conducted in the remaining ~106K participants (Stage 1). The top 69 SNPs associated with educational attainment were then carried forward for analysis in relation to cognitive performance in the ~24K subsample (Stage 2). In Stage 1, the chromosome 6q16.1 locus previously associated with years of schooling was again genome-wide significant for educational attainment, although the top SNP at this locus was slightly different (rs9320913 in the original GWAS and rs1487441 in the second GWAS; $R^2 > .9$ between the two SNPs). In Stage 2, rs1487441 was the SNP most strongly associated with cognitive performance ($P = 1.24 \times 10^{-4}$).

Notably, rs1487441 is a perfect proxy for our SNP, rs1906252 ($R^2 = 1$ in both HapMap and 1000 Genomes). Thus, we sought to integrate our current results with the most recent SSGAC findings. As shown in Table 3, we performed a meta-analysis of the SSGAC cognitive subcohort with our fully independent COGENT cohorts. LBC1936 and HBCS were part of the SSGAC studies, so they were excluded for this particular analysis. However, this was a conservative approach since (i) LBC1936 was included in the cognitive performance sample as part of the Childhood Intelligence Consortium, which used age 11 cognitive phenotypes for LBC1936, whereas LBC1936 adult phenotypes were included in COGENT; and (ii) HBCS was not included in the SSGAC cognitive analysis, only the

educational attainment GWAS.(Rietveld et al., 2014b) With a total sample size of 43,381 non-duplicated individuals across SSGAC and COGENT, the association between this locus and g had an estimated effect size of $\beta = .031$ ($P = 4.94 \times 10^{-7}$).

CHARGE Meta-Analysis

A GWAS of general cognitive function that included $\approx 53,000$ individuals from 31 population-based cohorts was recently published by the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium.(Davies et al., 2015) The study reported genome-wide significant SNP associations with general cognitive ability in three genomic regions, including rs1906252 ($P = 1.55 \times 10^{-8}$). We sought to integrate the COGENT findings with the CHARGE findings as well, and conducted a second meta-analysis of rs1906252 and g . In this analytic series, the COGENT samples were fully independent from both the SSGAC and the CHARGE samples. In other words, we excluded HBCS and LBC1936 as above, and excluded additional data from our Framingham Heart Study, Second Generation Cohort (FHS) sample, the Cardiovascular Health Study (CHS) sample, and the Norwegian Cognitive NeuroGenetics (NCNG) sample, all of which overlapped with CHARGE to some extent. In the reduced COGENT sample of 16 sites and $\approx 15,000$ subjects fully independent from SSGAC and CHARGE, rs1906252 retained a significant association to g ($\beta = .022$, $P = 2.81 \times 10^{-2}$). Further, the joint analysis of the 42 independent COGENT and CHARGE cohorts ($N \approx 68,000$) was robust ($\beta = .029$, $P = 1.65 \times 10^{-9}$).

Conclusions

The current study identified a significant association of a SNP at chromosome 6q16.1 with general cognitive ability across 21 well-characterized international samples of European-ancestry individuals from the general population. Sensitivity analysis suggested that the association between rs1906252 and g was significant, independent of the previously published cohorts from the educational attainment GWAS; moreover, the strength of the association was not affected by age of the subjects. In COGENT, the direction of association with g was consistent with the association to self-reported educational attainment in the SSGAC, with more copies of the minor A allele linked to better performance on objectively-measured sub-traits that comprise general cognitive ability.

Based on the commonly used formula,(Thorleifsson et al., 2009) $R^2 = 2f(1-f)a^2$, where f is the allele frequency (0.47 in this instance) and a is the additive effect as measured by standardized beta, the independent COGENT cohort produces an estimated effect size of $R^2 = .0382\%$ (Table 3). This effect size estimate is slightly more than half the value obtained in the SSGAC cognitive subcohort. This is probably due, at least in part, to the ‘winner’s curse’ phenomenon in which initial observations produce inflated effects by chance.(Zollner and Pritchard, 2007). The present report provides the first estimate of the effect size that is fully independent of the initial discovery cohort. This effect size is approximately an order of magnitude smaller than the largest allelic effect sizes observed for other complex polygenic traits such as height or weight (Visscher *et al*, 2012; Wood *et al*, 2014), possibly due to the inherent noise and heterogeneity in the construction of the g phenotype. However, the

success of GWAS for height and weight provide optimism that future, larger GWAS of cognitive ability will prove increasingly successful at identifying significant associations.

For example, the first study reporting a genome-wide significant QTL for height(Weedon et al., 2007) had power of only 3.2% to detect the *HMG2* locus, based on the effect size estimate later derived from an independent cohort that was an order of magnitude larger. (Lango Allen et al., 2010) Similarly, subsequent height GWAS(Weedon et al., 2008) remained underpowered by conventional criteria ($1-\beta = .80$) to detect even the strongest loci, yet were still able to obtain genome-wide significance for multiple loci, including those for which power was <1%. This pattern applies to all polygenic complex traits, due to the very large number of loci that are available to be detected, especially given that the number of associated loci tends to increase very rapidly as effect sizes drop.(Park et al., 2010)

Functionally, rs1906252 is about 700 kilobases from the nearest annotated gene, but it is in an intron of a long intergenic non-coding RNA (lincRNA; transcript ID: *RP11-436D23.1*; Gene ID: *LOC101927335*; Ensembl Gene ID: *ENSG00000271860*) that is expressed in human brain tissue based on searches in BrainSpan (<http://www.brainspan.org/>) and GeneProf(Halbritter et al., 2012) (<http://www.geneprof.org/>). In order to identify potential regulatory elements at this genomic locus, we interrogated rs1906252 using HaploReg v3 (Ward and Kellis, 2012). While rs1906252 is unannotated, it is in nearly perfect LD ($R^2 = .99$) with rs77910749, which is highly conserved (by both GERP and SiPhy computations), is a DNase hypersensitivity site in fetal brain, and serves as a weak enhancer or transcription start site across multiple brain tissues. Notably, rs1906252 was recently reported to be a genome-wide significant variant associated with bipolar disorder,(Mühleisen et al., 2014) again providing evidence of the important overlap of general cognitive ability and psychiatric illness; the primary finding of our previous COGENT study was a significant polygenic overlap between SNPs for general cognitive ability and schizophrenia.(Lencz et al., 2013) Additionally, rs1906252 was among the top findings (though not significant after correction for multiple comparisons) in an earlier GWAS of processing speed based on performance on the symbol search test conducted in part in the LBC1936 cohort.(Luciano et al., 2011) Most recently, rs1906252 and 10 other SNPs in this region were associated with general cognitive function in a GWAS of $\approx 53,000$ subjects at a threshold considered to reach genome-wide significance.(Davies et al., 2015)

In summary, the current study provides an independent replication of a link between genetic variation at rs1906252 and neurocognitive ability. Results also provide evidence that the proxy phenotype approach of using educational attainment as an indirect measure of cognitive ability for GWAS has external validity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Joey W. Trampush^{1,2,3,*}, Todd Lencz^{1,2,3}, Emma Knowles⁴, Gail Davies^{5,6}, Saurav Guha¹, Itsik Pe'er^{7,8}, David C. Liewald⁵, John M. Starr^{5,9}, Srdjan Djurovic^{10,11},

Ingrid Melle^{10,11,12}, Kjetil Sundet^{10,12}, Andrea Christoforou^{13,14}, Ivar Reinvang¹⁵, Semanti Mukherjee^{1,2}, Pamela DeRosse^{1,2}, Astri Lundervold^{16,17,18}, Vidar M. Steen^{13,14}, Majnu John^{1,2}, Thomas Espeseth^{15,19}, Katri Räikkönen²⁰, Elisabeth Widen²¹, Aarno Palotie^{21,22,23}, Johan G. Eriksson^{24,25,26,27,28}, Ina Giegling²⁹, Bettina Konte²⁹, Masashi Ikeda³⁰, Panos Roussos³¹, Stella Giakoumaki³², Katherine E. Burdick³¹, Antony Payton³³, William Ollier³³, Mike Horan³⁴, Matthew Scult³⁵, Dwight Dickinson³⁶, Richard E. Straub^{36,37}, Gary Donohoe³⁸, Derek Morris³⁸, Aiden Corvin³⁸, Michael Gill³⁸, Ahmad Hariri³⁵, Daniel R. Weinberger^{36,37}, Neil Pendleton³⁹, Nakao Iwata³⁰, Ariel Darvasi⁴⁰, Panos Bitsios⁴¹, Dan Rujescu²⁹, Jari Lahti^{20,42}, Stephanie Le Hellard^{13,14}, Matthew C. Keller⁴³, Ole A. Andreassen^{10,11,12}, Ian J. Deary^{5,6}, David C. Glahn⁴, and Anil K. Malhotra^{1,2,3}

Affiliations

¹Division of Psychiatry Research, Zucker Hillside Hospital, Glen Oaks, New York
²Center for Psychiatric Neuroscience, Feinstein Institute for Medical Research, Manhasset, New York
³Hofstra North Shore – LIJ School of Medicine, Department of Psychiatry, Hempstead, New York
⁴Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut
⁵Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, United Kingdom
⁶Department of Psychology, University of Edinburgh, Edinburgh, United Kingdom
⁷Department of Computer Science, Columbia University, New York, New York
⁸Center for Computational Biology and Bioinformatics, Columbia University, New York, New York
⁹Alzheimer Scotland Dementia Research Centre, University of Edinburgh, Edinburgh, United Kingdom
¹⁰NorMent, KG Jebsen Centre, Oslo, Norway
¹¹Oslo University Hospital, Oslo, Norway
¹²University of Oslo, Oslo, Norway
¹³K.G. Jebsen Centre for Psychosis Research, Dr. Einar Martens Research Group for Biological Psychiatry, Department of Clinical Medicine, University of Bergen, Bergen, Norway
¹⁴Center for Medical Genetics and Molecular Medicine, Haukeland University Hospital, Bergen, Norway
¹⁵Department of Psychology, University of Oslo, Oslo, Norway
¹⁶K.G. Jebsen Centre for Research on Neuropsychiatric Disorders, University of Bergen, Norway
¹⁷Department of Biological and Medical Psychology, University of Bergen, Norway
¹⁸Kavli Research Centre for Aging and Dementia, Haraldsplass Deaconess Hospital, Bergen, Norway
¹⁹K.G. Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway
²⁰Institute of Behavioural Sciences, University of Helsinki, Helsinki, Finland
²¹Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Finland
²²Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Cambridge, United Kingdom
²³Department of Medical Genetics, University of Helsinki and University Central Hospital, Helsinki, Finland
²⁴National Institute for Health and Welfare, Finland
²⁵Department of General Practice and Primary Health Care, University of Helsinki, Finland
²⁶Helsinki University Central Hospital, Unit of General Practice, Helsinki, Finland
²⁷Folkhälsan Research Centre, Helsinki, Finland
²⁸Vasa Central Hospital, Vasa, Finland
²⁹Department of Psychiatry, University of Halle, Halle, Germany
³⁰Department of Psychiatry, School of Medicine, Fujita Health University, Toyoake, Aichi, Japan
³¹Department of

Psychiatry, The Mount Sinai School of Medicine, New York, New York
³²Department of Psychology, School of Social Sciences, University of Crete, Greece
³³Centre for Integrated Genomic Medical Research, University of Manchester, Manchester, United Kingdom ³⁴School of Community-Based Medicine, Neurodegeneration Research Group, University of Manchester, Manchester, United Kingdom ³⁵Laboratory of NeuroGenetics, Department of Psychology & Neuroscience, Duke University, Durham, North Carolina ³⁶Clinical Brain Disorders Brain and Genes, Cognition and Psychosis Program, Intramural Research Program, National Institute of Mental Health, National Institute of Health, Bethesda, Maryland ³⁷Lieber Institute for Brain Development, Johns Hopkins University Medical Campus, Baltimore, Maryland ³⁸Neuropsychiatric Genetics Research Group, Department of Psychiatry and Trinity College Institute of Neuroscience, Trinity College Dublin, Dublin, Ireland ³⁹Institute of Brain, Behaviour and Mental Health, University of Manchester, Manchester, United Kingdom ⁴⁰Department of Genetics, The Institute of Life Sciences, The Hebrew University of Jerusalem, Jerusalem, Israel ⁴¹Department of Psychiatry and Behavioral Sciences, Faculty of Medicine, University of Crete, Heraklion, Crete, Greece ⁴²Institute of Genetics, Folkhälsan Research Centre, Helsinki, Finland ⁴³Institute for Behavioral Genetics, University of Colorado, Boulder, Colorado

Acknowledgements

This work has been supported by grants from the National Institutes of Health (R01 MH079800 and P50 MH080173 to AKM; RC2 MH089964 to TL; R01 MH080912 to DCG; K23 MH077807 to KEB; K01 MH085812 to MCK). Dr. Donohoe is generously funded by the Health Research Board (Ireland) and Science Foundation Ireland. Data collection for the TOP cohort was supported by the Research Council of Norway, South-East Norway Health Authority. The NCNG study was supported by Research Council of Norway Grants 154313/V50 and 177458/V50. The NCNG GWAS was financed by grants from the Bergen Research Foundation, the University of Bergen, the Research Council of Norway (FUGE, Psykisk Helse), Helse Vest RHF and Dr. Einar Martens Fund. The Helsinki Birth Cohort Study has been supported by grants from the Academy of Finland, the Finnish Diabetes Research Society, Folkhälsan Research Foundation, Novo Nordisk Foundation, Finska Läkaresällskapet, Signe and Ane Gyllenberg Foundation, University of Helsinki, Ministry of Education, Ahokas Foundation, Emil Aaltonen Foundation. For the LBC1936 cohort we thank the cohort participants and team members who contributed to this study. Phenotype collection was supported by Age UK (The Disconnected Mind project). Genotyping was funded by the UK Biotechnology and Biological Sciences Research Council (BBSRC). The work was undertaken by The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology, part of the cross council Lifelong Health and Wellbeing Initiative (MR/K026992/1). Funding from the BBSRC and Medical Research Council (MRC) is gratefully acknowledged. The Duke Neurogenetic Study was supported by an NSF Graduate Research Fellowship to MS, Duke University, and NIDA grants R01DA031579 & R01DA033369. The NIMH Genes, Cognition and Psychosis Study was funded by the NIH/NIMH Intramural Research Program and the Lieber Institute for Brain Development. Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by

the Laboratory for Neuro Imaging at the University of Southern California. Finally, several publicly available datasets were included; we kindly thank the investigative teams and staffs of the Pediatric Imaging, Neurocognition, and Genetics (PING) study, the Alzheimer's Disease Neuroimaging Initiative (ADNI) project, and the studies who made their data available in dbGaP.

References

- Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. GenABEL: an R library for genome-wide association analysis. *Bioinformatics*. 2007; 23:1294–6. [PubMed: 17384015]
- Batty GD, Shipley MJ, Gale CR, Mortensen LH, Deary IJ. Does IQ predict total and cardiovascular disease mortality as strongly as other risk factors? Comparison of effect estimates using the Vietnam Experience Study. *Heart*. 2008; 94:1541–4. [PubMed: 18801778]
- Benyamin B, Pourcain B, Davis OS, Davies G, Hansell NK, Brion M-J a, Kirkpatrick RM, Cents R a M, Frani S, Miller MB, Haworth CM a, Meaburn E, Price TS, Evans DM, Timpson N, Kemp J, Ring S, McArdle W, Medland SE, Yang J, Harris SE, Liewald DC, Scheet P, Xiao X, Hudziak JJ, de Geus EJC, Jaddoe VWV, Starr JM, Verhulst FC, Pennell C, Tiemeier H, Iacono WG, Palmer LJ, Montgomery GW, Martin NG, Boomsma DI, Posthuma D, McGue M, Wright MJ, Davey Smith G, Deary IJ, Plomin R, Visscher PM. Childhood intelligence is heritable, highly polygenic and associated with FBNP1L. *Mol Psychiatry*. 2014; 19:253–8. [PubMed: 23358156]
- Carroll, JB. *Human Cognitive Abilities: A Survey of Factor-Analytic Studies*. Cambridge University Press; Cambridge, UK: 1993.
- Chabris CF, Hebert BM, Benjamin DJ, Beauchamp J, Cesarini D, van der Loos M, Johannesson M, Magnusson PKE, Lichtenstein P, Atwood CS, Freese J, Hauser TS, Hauser RM, Christakis N, Laibson D. Most reported genetic associations with general intelligence are probably false positives. *Psychol Sci*. 2012; 23:1314–23. [PubMed: 23012269]
- Chen J, Lipska BK, Halim N, Ma QD, Matsumoto M, Melhem S, Kolachana BS, Hyde TM, Herman MM, Apud J, Egan MF, Kleinman JE, Weinberger DR. Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in postmortem human brain. *Am J Hum Genet*. 2004; 75:807–821. [PubMed: 15457404]
- Davies G, Armstrong N, Bis JC, Bressler J, Chouraki V, Giddaluru S, Hofer E, Ibrahim verbaas C a, Kirin M, Lahti J. Genetic contributions to variation in general cognitive function: a meta-analysis of genome-wide association studies in the CHARGE consortium (N = 53 949). *Mol Psychiatry*. 2015; 20(2):183–92. [PubMed: 25644384]
- Davies G, Tenesa A, Payton A, Yang J, Harris SE, Liewald D, Ke X, Le Hellard S, Christoforou A, Luciano M, McGhee K, Lopez L, Gow AJ, Corley J, Redmond P, Fox HC, Haggarty P, Whalley LJ, McNeill G, Goddard ME, Espeseth T, Lundervold a J, Reinvang I, Pickles A, Steen VM, Ollier W, Porteous DJ, Horan M, Starr JM, Pendleton N, Visscher PM, Deary IJ. Genome-wide association studies establish that human intelligence is highly heritable and polygenic. *Mol Psychiatry*. 2011; 16:996–1005. [PubMed: 21826061]
- Deary IJ. Intelligence. *Annu Rev Psychol*. 2012; 63:453–82. [PubMed: 21943169]
- Deary IJ. Why do intelligent people live longer? *Nature*. 2008; 456:175–76. [PubMed: 19005537]
- Deary IJ, Johnson W, Houlihan LM. Genetic foundations of human intelligence. *Hum Genet*. 2009; 126:215–32. [PubMed: 19294424]
- Deary IJ, Penke L, Johnson W. The neuroscience of human intelligence differences. *Nat Rev Neurosci*. 2010; 11:201–211. [PubMed: 20145623]
- Deary IJ, Weiss A, Batty GD. Intelligence and personality as predictors of illness and death: how researchers in differential psychology and chronic disease epidemiology are collaborating to understand and address health inequalities. *Psychol Sci Public Interes*. 2011; 11:53–79.
- Deary IJ, Yang J, Davies G, Harris SE, Tenesa A, Liewald D, Luciano M, Lopez LM, Gow AJ, Corley J, Redmond P, Fox HC, Rowe SJ, Haggarty P, McNeill G, Goddard ME, Porteous DJ, Whalley LJ, Starr JM, Visscher PM. Genetic contributions to stability and change in intelligence from childhood to old age. *Nature*. 2012; 482:212–5. [PubMed: 22258510]
- Donohoe G, Deary IJ, Glahn DC, Malhotra AK, Burdick KE. Neurocognitive phenomics: examining the genetic basis of cognitive abilities. *Psychol Med*. 2013; 43:2027–361–10. [PubMed: 23194477]

- Gottfredson LS. Why g matters: The complexity of everyday life. *Intelligence*. 1997; 24:79–132.
- Halbritter F, Vaidya HJ, Tomlinson SR. GeneProf: analysis of high-throughput sequencing experiments. *Nat Meth*. 2012; 9:7–8.
- Han B, Eskin E. Random-effects model aimed at discovering associations in meta-analysis of genome-wide association studies. *Am J Hum Genet*. 2011; 88:586–98. [PubMed: 21565292]
- Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet*. 2009; 5:e1000529. [PubMed: 19543373]
- Johnson W, Deary IJ, Iacono WG. Genetic and environmental transactions underlying educational attainment. *Intelligence*. 2009; 37:466–478. [PubMed: 20161120]
- Johnson W, te Nijenhuis J, Bouchard TJ Jr. Still just 1 g: Consistent results from five test batteries. *Intelligence*. 2008; 36:81–95.
- Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F, Willer CJ, Jackson AU, Vedantam S, Raychaudhuri S, Ferreira T, Wood AR, Weyant RJ, Segrè AV, Speliotes EK, Wheeler E, Soranzo N, Park J-H, Yang J, Gudbjartsson D, Heard-Costa NL, Randall JC, Qi L, Vernon Smith A, Mägi R, Pastinen T, Liang L, Heid IM, Luan J, Thorleifsson G, Winkler TW, Goddard ME, Sin Lo K, Palmer C, Workalemahu T, Aulchenko YS, Johansson A, Zillikens MC, Feitosa MF, Esko T, Johnson T, Ketkar S, Kraft P, Mangino M, Prokopenko I, Absher D, Albrecht E, Ernst F, Glazer NL, Hayward C, Hottenga J-J, Jacobs KB, Knowles JW, Kutalik Z, Monda KL, Polasek O, Preuss M, Rayner NW, Robertson NR, Steinthorsdottir V, Tyrer JP, Voight BF, Wiklund F, Xu J, Zhao JH, Nyholt DR, Pellikka N, Perola M, Perry JRB, Surakka I, Tammesoo M-L, Altmaier EL, Amin N, Aspelund T, Bhangale T, Boucher G, Chasman DI, Chen C, Coin L, Cooper MN, Dixon AL, Gibson Q, Grundberg E, Hao K, Juhani Juntila M, Kaplan LM, Kettunen J, König IR, Kwan T, Lawrence RW, Levinson DF, Lorentzon M, McKnight B, Morris AP, Müller M, Suh Ngwa J, Purcell S, Rafelt S, et al. Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature*. 2010; 467:832–8. [PubMed: 20881960]
- Lencz T, Knowles E, Davies G, Guha S, Liewald DC, Starr JM, Djurovic S, Melle I, Sundet K, Christoforou A, Reinvang I, Mukherjee S, Derosse P, Lundervold A, Steen VM, John M, Espeseth T, Rääkkönen K, Widen E, Palotie A, Eriksson JG, Giegling I, Konte B, Ikeda M, Roussos P, Giakoumaki S, Burdick KE, Payton A, Ollier W, Horan M, Donohoe G, Morris D, Corvin A, Gill M, Pendleton N, Iwata N, Darvasi A, Bitsios P, Rujescu D, Lahti J, Hellard SL, Keller MC, Andreassen OA, Deary IJ, Glahn DC, Malhotra AK. Molecular genetic evidence for overlap between general cognitive ability and risk for schizophrenia: a report from the Cognitive Genomics consortium (COGENT). *Mol Psychiatry*. 2014; 19:168–74. [PubMed: 24342994]
- Luciano M, Hansell NK, Lahti J, Davies G, Medland SE, Rääkkönen K, Tenesa A, Widen E, McGhee KA, Palotie A, Liewald D, Porteous DJ, Starr JM, Montgomery GW, Martin NG, Eriksson JG, Wright MJ, Deary IJ. Whole genome association scan for genetic polymorphisms influencing information processing speed. *Biol Psychol*. 2011; 86:193–202. [PubMed: 21130836]
- Marioni RE, Davies G, Hayward C, Liewald D, Kerr SM, Campbell A, Luciano M, Smith BH, Padmanabhan S, Hocking LJ, Hastie ND, Wright AF, Porteous DJ, Visscher PM, Deary IJ. Molecular genetic contributions to socioeconomic status and intelligence. *Intelligence*. 2014; 44:26–32. [PubMed: 24944428]
- Martin NW, Medland SE, Verweij KJH, Lee SH, Nyholt DR, Madden P a, Heath AC, Montgomery GW, Wright MJ, Martin NG. Educational attainment: a genome wide association study in 9538 Australians. *PLoS One*. 2011; 6:e20128. [PubMed: 21694764]
- Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS, Connor R, Davis S, Deakin B, DeRubeis RJ, Dubois B, Geyer M a, Goodwin GM, Gorwood P, Jay TM, Joëls M, Mansuy IM, Meyer-Lindenberg A, Murphy D, Rolls E, Saletu B, Spedding M, Sweeney J, Whittington M, Young LJ. Cognitive dysfunction in psychiatric disorders: characteristics, causes and the quest for improved therapy. *Nat Rev Drug Discov*. 2012; 11:141–68. [PubMed: 22293568]
- Mühleisen TW, Leber M, Schulze TG, Strohmaier J, Degenhardt F, Treutlein J, Mattheisen M, Forstner AJ, Schumacher J, Breuer R, Meier S, Herms S, Hoffmann P, Lacour A, Witt SH, Reif A, Müller-Myhsok B, Lucae S, Maier W, Schwarz M, Vedder H, Kammerer-Ciernioch J, Pfennig A, Bauer M, Hautzinger M, Moebus S, Priebe L, Czerski PM, Hauser J, Lissowska J, Szeszenia-Dabrowska N, Brennan P, McKay JD, Wright A, Mitchell PB, Fullerton JM, Schofield PR,

- Montgomery GW, Medland SE, Gordon SD, Martin NG, Krasnow V, Chuchalin A, Babadjanova G, Pantelejeva G, Abramova LI, Tiganov AS, Polonikov A, Khusnutdinova E, Alda M, Grof P, Rouleau G a, Turecki G, Laprise C, Rivas F, Mayoral F, Kogevinas M, Grigoriou-Serbanescu M, Propping P, Becker T, Rietschel M, Nöthen MM, Cichon S. Genome-wide association study reveals two new risk loci for bipolar disorder. *Nat Commun.* 2014; 5:3339. [PubMed: 24618891]
- Need AC, Attix DK, McEvoy JM, Cirulli ET, Linney KL, Hunt P, Ge D, Heinzen EL, Maia JM, Shianna KV, Weale ME, Cherkas LF, Clement G, Spector TD, Gibson G, Goldstein DB. A genome-wide study of common SNPs and CNVs in cognitive performance in the CANTAB. *Hum Mol Genet.* 2009; 18:4650–61. [PubMed: 19734545]
- Panizzon MS, Vuoksima E, Spoon KM, Jacobson KC, Lyons MJ, Franz CE, Xian H, Vasilopoulos T, Kremen WS. Genetic and environmental influences on general cognitive ability: Is g a valid latent construct? *Intelligence.* 2014; 43:65–76. [PubMed: 24791031]
- Park J-H, Wacholder S, Gail MH, Peters U, Jacobs KB, Chanock SJ, Chatterjee N. Estimation of effect size distribution from genome-wide association studies and implications for future discoveries. *Nat Genet.* 2010; 42:570–5. [PubMed: 20562874]
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007; 81:559–75. [PubMed: 17701901]
- Rietveld CA, Conley D, Eriksson N, Esko T, Medland SE, Vinkhuyzen AAE, Yang J, Boardman JD, Chabris CF, Dawes CT, Domingue BW, Hinds DA, Johannesson M, Kiefer AK, Laibson D, Magnusson PKE, Mountain JL, Oskarsson S, Rostapshova O, Teumer A, Tung JY, Visscher PM, Benjamin DJ, Cesarini D, Koellinger PD, Consortium, the SSGA. Replicability and robustness of genome-wide association studies for behavioral traits. *Psychol Sci.* 2014a; 25:1975–86. [PubMed: 25287667]
- Rietveld CA, Esko T, Davies G, Pers TH, Turley P, Benyamin B, Chabris CF, Emilsson V, Johnson AD, Lee JJ, de Leeuw C, Marioni RE, Medland SE, Miller MB, Rostapshova O, van der Lee SJ, Vinkhuyzen AAE, Amin N, Conley D, Derringer J, van Duijn CM, Fehrmann R, Franke L, Glaeser EL, Hansell NK, Hayward C, Iacono WG, Ibrahim-Verbaas C, Jaddoe V, Karjalainen J, Laibson D, Lichtenstein P, Liewald DC, Magnusson PKE, Martin NG, McGue M, McMahon G, Pedersen NL, Pinker S, Porteous DJ, Posthuma D, Rivadeneira F, Smith BH, Starr JM, Tiemeier H, Timpson NJ, Trzaskowski M, Uitterlinden AG, Verhulst FC, Ward ME, Wright MJ, Davey Smith G, Deary IJ, Johannesson M, Plomin R, Visscher PM, Benjamin DJ, Cesarini D, Koellinger PD. Common genetic variants associated with cognitive performance identified using the proxy-phenotype method. *Proc Natl Acad Sci U S A.* 2014b; 111:13790–4. [PubMed: 25201988]
- Rietveld CA, Medland SE, Derringer J, Yang J, Esko T, Martin NG, Westra H-J, Shakhbazov K, Abdellaoui A, Agrawal A, Albrecht E, Alizadeh BZ, Amin N, Barnard J, Baumeister SE, Benke KS, Bielak LF, Boatman J a, Boyle P a, Davies G, de Leeuw C, Eklund N, Evans DSDM, Fehrmann R, Fischer K, Gieger C, Gjessing HK, Hägg S, Harris JRJM, Hayward C, Holzappel C, Ibrahim-Verbaas C a, Ingelsson E, Jacobsson B, Joshi PK, Jugessur A, Kaakinen M, Kanoni S, Karjalainen J, Kolcic I, Kristiansson K, Kutalik Z, Lahti J, Lee SH, Lin P, Lind P a, Liu Y, Lohman K, Loitfelder M, McMahon G, Vidal PM, Meirelles O, Milani L, Myhre R, Nuotio M-L, Oldmeadow CJ, Petrovic KE, Peyrot WJ, Polasek O, Quaye L, Reinmaa E, Rice JP, Rizzi TS, Schmidt H, Schmidt R, Smith AV, Smith J a, Tanaka T, Terracciano A, van der Loos MJHM, Vitart V, Völzke H, Wellmann J, Yu L, Zhao W, Allik J, Attia JR, Bandinelli S, Bastardot F, Beauchamp J, Bennett D a, Berger K, Bierut LJ, Boomsma DI, Bültmann U, Campbell H, Chabris CF, Cherkas L, Chung MK, Cucca F, de Andrade M, De Jager PL, De Neve J-E, Deary IJ, Dedoussis GV, Deloukas P, Dimitriou M, Eiriksdóttir G, et al. GWAS of 126,559 individuals identifies genetic variants associated with educational attainment. *Science (80-).* 2013; 340:1467–71.
- Thorleifsson G, Walters GB, Gudbjartsson DF, Steinthorsdóttir V, Sulem P, Helgadóttir A, Styrkarsdóttir U, Gretarsdóttir S, Thorlacius S, Jonsdóttir I, Jonsdóttir T, Olafsdóttir EJ, Olafsdóttir GH, Jonsson T, Jonsson F, Borch-Johnsen K, Hansen T, Andersen G, Jorgensen T, Lauritzen T, Aben KK, Verbeek ALM, Roeleveld N, Kampman E, Yanek LR, Becker LC, Tryggvadóttir L, Rafnar T, Becker DM, Gulcher J, Kiemeny L a, Pedersen O, Kong A, Thorsteinsdóttir U, Stefansson K. Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet.* 2009; 41:18–24. [PubMed: 19079260]

- Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res.* 2012; 40:D930–4. [PubMed: 22064851]
- Weedon MN, Lango H, Lindgren CM, Wallace C, Evans DM, Mangino M, Freathy RM, Perry JRB, Stevens S, Hall AS, Samani NJ, Shields B, Prokopenko I, Farrall M, Dominiczak A, Johnson T, Bergmann S, Beckmann JS, Vollenweider P, Waterworth DM, Mooser V, Palmer CN a, Morris AD, Ouwehand WH, Zhao JH, Li S, Loos RJF, Barroso I, Deloukas P, Sandhu MS, Wheeler E, Soranzo N, Inouye M, Wareham NJ, Caulfield M, Munroe PB, Hattersley AT, McCarthy MI, Frayling TM. Genome-wide association analysis identifies 20 loci that influence adult height. *Nat Genet.* 2008; 40:575–83. [PubMed: 18391952]
- Weedon MN, Lettre G, Freathy RM, Lindgren CM, Voight BF, Perry JRB, Elliott KS, Hackett R, Guiducci C, Shields B, Zeggini E, Lango H, Lyssenko V, Timpson NJ, Burt NP, Rayner NW, Saxena R, Ardlie K, Tobias JH, Ness AR, Ring SM, Palmer CN a, Morris AD, Peltonen L, Salomaa V, Davey Smith G, Groop LC, Hattersley AT, McCarthy MI, Hirschhorn JN, Frayling TM. A common variant of HMGA2 is associated with adult and childhood height in the general population. *Nat Genet.* 2007; 39:1245–50. [PubMed: 17767157]
- Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 2011; 88:76–82. [PubMed: 21167468]
- Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the application of mixed-model association methods. *Nat Genet.* 2014; 46:100–6. [PubMed: 24473328]
- Zollner S, Pritchard JK. Overcoming the winner's curse: estimating penetrance parameters from case-control data. *Am J Hum Genet.* 2007; 80:605–15. [PubMed: 17357068]




Figure 1.

Kernel density plot (KDP) of the g factor across 21 COGENT studies. Each g score is smoothed for each individual using a strongly peaked kernel function in order to evaluate every point along the x -axis. The y -axis represents density. As can be seen, the shape of the distribution tightly fits a Gaussian curve across all studies.




Figure 2. Results (in forest-plot format) of the meta-analysis between rs1906252 and general cognitive ability in all 21 COGENT studies. A positive effect was detected in 17 out of 21 studies.




Figure 3. Effect plot of the association between rs1906252 and general cognitive ability. A linear increase in general cognitive ability was related positively to minor A allele load. In the combined sample, 5,254 subjects had no copies of the minor allele, 10,329 subjects were heterozygous, and 4,912 subjects had two copies of the minor A allele.

Table 1

Sites and basic demographics of samples included in the Cognitive Genomics Consortium (COGENT; ordered alphabetically).

Study	Study Name	N	Age (mean)	Age range	% Male
ADNI1	Alzheimer's Disease Neuroimaging Initiative, Phase 1	127	75.7	62-90	55.9
CHS	Cardiovascular Health Study	2,114	77.9	69-99	37.1
DNS	Duke Neurogenetics Study	357	19.7	18-22	47.6
FHS	Framingham Heart Study, Second Generation Cohort	1,650	65.3	38-90	45.6
GCAP	NIMH Genes, Cognition and Psychosis Program	655	31.5	18-60	46.9
GenADA	Genotype-Phenotype Associations in Alzheimer's Disease	782	73.4	48-94	35.7
HBCS	Helsinki Birth Cohort Study	332	67.7	64-75	100.0
IBG	Institute for Behavioral Genetics	299	15.9	12-19	78.6
LBC1936	Lothian Birth Cohort 1936	1,005	69.6	68-71	50.6
LLFS	Long Life Family Study	3,606	64.6	24-89	45.2
LOAD	Late Onset Alzheimer's Disease Family Study	1,141	75.1	53-98	36.6
LOGOS	Learning on Genetics of Schizophrenia Spectrum	864	22.5	18-44	100.0
MAN	Manchester Longitudinal Studies of Cognitive Aging	805	67.2	45-87	28.9
MUC1	Munich, Germany Sample 1	588	50.5	19-79	50.0
MUC2	Munich, Germany Sample 2	538	45.1	19-72	45.7
NCNG	Norwegian Cognitive NeuroGenetics	670	47.6	18-79	31.9
NEW	Newcastle Longitudinal Studies of Cognitive Aging	753	67.0	54-87	29.0
PING	Pediatric Imaging, Neurocognition and Genetics Study	637	11.6	3-18	52.4
PNC	Philadelphia Neurodevelopmental Cohort	4,412	13.8	8-21	50.2
TOP	Thematic Organized Psychosis Research Study	394	34.6	17-55	49.2
ZHH	Zucker Hillside Hospital	219	35.3	8-78	49.3

Table 2

Results of the association between rs1906252 and general cognitive ability in COGENT.

Data set	Studies (N)	Subjects (N)	Fixed effects			Random effects			Heterogeneity estimates			
			β	S.E.	P	β	S.E.	P	I ²	Q	PQ	Tau ²
COGENT1	9	4,962	.050	.020	1.20E-02	.050	.022	2.08E-02	11.411	9.030	.340	0
COGENT2	12	15,533	.027	.009	2.34E-03	.027	.009	2.34E-03	0	8.472	.671	0
COGENT, all cohorts	21	20,495	.031	.008	1.47E-04	.031	.008	1.47E-04	0	18.668	.543	0
COGENT, excluding HBGS and LBC1936 ^d	19	19,192	.026	.008	1.60E-03	.026	.008	1.60E-03	0	11.126	.889	0
COGENT, excluding HBGS, LBC1936, CHS, FHS and NCNG ^b	16	14,971	.022	.010	2.81E-02	.022	.010	2.81E-02	0	10.467	.789	0

Abbreviations: COGENT, Cognitive Genomics Consortium; COGENT1, nine sites included in COGENT Phase 1 (Lencz et al, 2014); COGENT2, 12 sites added to Phase 2 of COGENT; HBGS, Helsinki Birth Cohort Study; LBC1936, Lothian Birth Cohort 1936; CHS, Cardiovascular Health Study; FHS, Framingham Heart Study; NCNG, Norwegian Cognitive NeuroGenetics Study

^a HBGS and LBC1936 removed since both studies were part of the Social Science Genetic Association Consortium (SSGAC) educational attainment GWAS study (Rietveld et al, 2013) and/or SSGAC cognition study (Rietveld et al., 2014).

^b HBGS and LBC1936 removed since both studies were part of the SSGAC educational attainment GWAS study (Rietveld et al, 2013) and/or SSGAC cognition study (Rietveld et al., 2014), and CHS, FHS and NCNG removed since these studies were part of the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium cognition study (Davies et al, 2015).

Table 3

Meta-analytic results of the association between general cognitive ability and genetic variants in chromosome 6q16.1 associated with educational attainment in COGENT (SNP rs1906252), SSGAC (SNP rs1487441) and CHARGE (SNP rs1906252).

Consortium	<i>N</i> studies	<i>N</i> subjects	β	S.E.	<i>P</i>
COGENT, excluding HBCS and LBC1936	19	19,192	.026	.008	1.60E-03
SSGAC	11	24,189	.036	.009	1.24E-04
COGENT and SSGAC	30	43,381	.031	.006	4.94E-07
COGENT, excluding HBCS, LBC1936, CHS, FHS and NCNG	16	14,971	.022	.010	2.81E-02
CHARGE	26	53,188	.031	.006	1.55E-08
COGENT and CHARGE	42	68,159	.029	.005	1.65E-09

Abbreviations: COGENT, Cognitive Genomics Consortium; SSGAC, Social Science Genetic Association Consortium; CHARGE, Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium. *Note:* COGENT studies that overlapped with SSGAC (HBCS and LBC1936) and/or CHARGE (HBCS, LBC1936, CHS, FHS, and NCNG) were removed prior to conducting joint analyses. The Lothian Birth Cohort Studies of 1921 and 1936 and the Helsinki Birth Cohort Study (HBCS) overlapped between SSGAC and CHARGE.