

2014

Prognostic and biologic significance of DNMT3B expression in older patients with cytogenetically normal primary acute myeloid leukemia

C. Niederwieser

J. Kohlschmidt

S. Volinia

S. P. Whitman

K. H. Metzeler

See next page for additional authors

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>



Part of the [Hematology Commons](#), and the [Oncology Commons](#)

Recommended Citation

Niederwieser C, Kohlschmidt J, Volinia S, Whitman S, Metzeler K, Eisfeld A, Maharry K, Yan P, Kolitz J, Bloomfield C, . Prognostic and biologic significance of DNMT3B expression in older patients with cytogenetically normal primary acute myeloid leukemia. . 2014 Jan 01; 29(3):Article 2432 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/2432>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

Authors

C. Niederwieser, J. Kohlschmidt, S. Volinia, S. P. Whitman, K. H. Metzeler, A. K. Einfeld, K. Maharry, P. Yan, J. E. Kowitz, C. D. Bloomfield, and +17 additional authors



Published in final edited form as:

Leukemia. 2015 March ; 29(3): 567–575. doi:10.1038/leu.2014.267.

Prognostic and biologic significance of *DNMT3B* expression in older patients with cytogenetically normal primary acute myeloid leukemia

Christian Niederwieser¹, Jessica Kohlschmidt^{1,2}, Stefano Volinia¹, Susan P. Whitman¹, Klaus H. Metzeler¹, Ann-Kathrin Eisfeld¹, Kati Maharry^{1,2}, Pearly Yan¹, David Frankhouser¹, Heiko Becker¹, Sebastian Schwind¹, Andrew J. Carroll³, Deedra Nicolet^{1,2}, Jason H. Mendler¹, John P. Curfman¹, Yue-Zhong Wu¹, Maria R. Baer⁴, Bayard L. Powell⁵, Jonathan E. Kolitz⁶, Joseph O. Moore⁷, Thomas H. Carter⁸, Ralf Bundschuh⁹, Richard A. Larson¹⁰, Richard M. Stone¹¹, Krzysztof Mrózek^{1,*}, Guido Marcucci^{1,*}, and Clara D. Bloomfield^{1,*}

¹The Ohio State University Comprehensive Cancer Center, Columbus, OH

²Alliance for Clinical Trials in Oncology Statistics and Data Center, Mayo Clinic, Rochester, MN

³University of Alabama at Birmingham, Birmingham, AL

⁴Greenebaum Cancer Center, University of Maryland, Baltimore, MD

⁵Comprehensive Cancer Center of Wake Forest University, Winston-Salem, NC

⁶Monter Cancer Center, Hofstra North Shore-Long Island Jewish School of Medicine, Lake Success, NY

⁷Duke University Medical Center, Durham, NC

⁸University of Iowa, Iowa City, IA

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence: Clara D. Bloomfield, MD, The Ohio State University Comprehensive Cancer Center, 1216 James Cancer Hospital, 300 West Tenth Avenue, Columbus, OH 43210-1228. clara.bloomfield@osumc.edu; or Guido Marcucci, MD, The Ohio State University Comprehensive Cancer Center, 410 Biomedical Research Tower, 460 West Twelfth Avenue, Columbus, OH 43210. guido.marcucci@osumc.edu; or Krzysztof Mrózek, MD, PhD, The Ohio State University Comprehensive Cancer Center, 1232A James Cancer Hospital, 300 West Tenth Avenue, Columbus, OH 43210-1228. krzysztof.mrozek@osumc.edu.

*These senior authors contributed equally to this work.

Presented in part at the 18th Congress of the European Hematology Association, Stockholm, Sweden, June 13–16, 2013, and published in abstract form: Niederwieser C, Kohlschmidt J, Maharry K, Mrózek K, Metzeler K, Volinia S *et al*. High expression of DNMT3B negatively impacts on clinical outcome of older patients (pts) with primary cytogenetically normal (CN) acute myeloid leukemia (AML) [CALGB 20202 (ALLIANCE)]. *Haematologica* 2013; **98**(suppl 1): 484 (abstract S1168).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

CN, JK, SV, K Mrózek, GM and CDB designed the study, analyzed the data and wrote the manuscript, and all authors agreed on the final version; SPW, KHM, A-KE, PY, DF HB, SS, JHM, JPC, Y-ZW, and RB carried out laboratory-based research; JK, K Maharry, SV, and DN performed statistical analyses; and AJC, MRB, BLP, JEK, JOM, THC, RAL, RMS, K Mrózek, GM and CDB were involved directly or indirectly in the care of patients and/or sample procurement.

Supplementary Information is available at *Leukemia's* website (<http://www.nature.com/leu>)

⁹Departments of Physics and Chemistry & Biochemistry, The Ohio State University, Columbus, OH

¹⁰University of Chicago Medical Center, Chicago, IL

¹¹Dana Farber Cancer Institute, Boston, MA

Abstract

DNMT3B encodes a DNA methyltransferase implicated in aberrant epigenetic changes contributing to leukemogenesis. We tested whether *DNMT3B* expression, measured by NanoString nCounter assay, associates with outcome, gene- and microRNA-expression and DNA methylation profiles in 210 older (> 60 years) adults with primary, cytogenetically normal AML (CN-AML). Patients were dichotomized into high versus low expressers using median cut. Outcomes were assessed in the context of known CN-AML prognosticators. Gene- and microRNA-expression, and DNA methylation profiles were analyzed using microarrays and MethylCap-sequencing, respectively. High *DNMT3B* expressers had fewer complete remissions (CR; $P=0.002$) and shorter disease-free (DFS; $P=0.02$) and overall (OS; $P<0.001$) survival. In multivariable analyses, high *DNMT3B* expression remained an independent predictor of lower CR rates ($P=0.04$) and shorter DFS ($P=0.04$) and OS ($P=0.001$). High *DNMT3B* expression associated with a gene-expression profile comprising 363 genes involved in differentiation, proliferation and survival pathways, but with only 4 differentially expressed microRNAs (*miR-133b*, *miR-148a*, *miR-122*, *miR-409-3p*) and no differential DNA methylation regions. We conclude that high *DNMT3B* expression independently associates with adverse outcome in older CN-AML patients. Gene-expression analyses suggest that *DNMT3B* is involved in the modulation of several genes, although the regulatory mechanisms remain to be investigated to devise therapeutic approaches specific for these patients.

Keywords

acute myeloid leukemia; *DNMT3B* expression; prognostication; gene-expression profiling

INTRODUCTION

Acute myeloid leukemia (AML) is a heterogeneous disease presenting with a wide spectrum of prognostically relevant cytogenetic aberrations, gene mutations and abnormal expression of genes and microRNAs. Cytogenetically normal AML (CN-AML) patients, constituting 40 to 50% of all AML patients,¹ are the largest and molecularly best characterized cytogenetic subset in primary (*de novo*) AML.¹⁻³ Although leukemic blasts of these patients do not contain microscopically detectable chromosome abnormalities, they harbor prognostically relevant mutations and aberrantly expressed genes and microRNAs.²⁻¹⁶ In addition to these genetic alterations, epigenetic changes have recently been shown to participate in myeloid leukemogenesis and be pharmacologically targetable.^{17,18} Notably, some genes whose mutations are prognostic in CN-AML encode proteins that are implicated in epigenetic regulation of gene transcription, namely *IDH2*, *ASXL1* and *DNMT3A*. The latter is among the most frequently mutated genes in primary CN-AML patients, being found mutated in 29 to 34% of the patients.^{9,19}

DNMT3A encodes DNA methyltransferase 3A (DNMT3A), which is involved in epigenetic gene silencing through DNA hypermethylation.²⁰ In addition to DNMT3A, DNMT1 and DNMT3B also mediate DNA methylation in normal and malignant cells, and may represent potential therapeutic targets in cancer and leukemia.^{21–24} However, in contrast to *DNMT3A*, no recurrent mutations of *DNMT1* and *DNMT3B* genes have been reported in AML.²⁵ Instead, one study has indicated that higher expression of *DNMT3B* is associated with worse outcome in AML.²⁶ However, the patient cohort analyzed was cytogenetically diverse and heterogeneous for clinical features and treatment received. Thus, it is unknown whether *DNMT3B* expression is an independent prognostic factor and can be used for stratification guidance in CN-AML.

Thus, we analyzed the clinical significance of *DNMT3B* expression in the context of a comprehensive panel of molecular prognosticators in a relatively large cohort of older (aged 60 years) patients with CN-AML who were similarly treated on cytarabine/daunorubicin-based protocols. To gain biologic insights, we also derived genome-wide *DNMT3B*-associated gene- and microRNA-expression and DNA methylation profiles. We studied older patients because both the incidence of AML and the role of epigenetics increase with age. Moreover, we have recently reported a favorable clinical response to hypomethylating agents in this age group of AML patients.²⁷

PATIENTS AND METHODS

Patients, treatment and cytogenetic studies

Pretreatment bone marrow (BM) or blood samples were obtained from 210 patients with primary CN-AML aged 60 to 83 years (median, 68 years) who received intensive first-line therapy on Cancer and Leukemia Group B (CALGB) trials.^{28–32} All patients received cytarabine-daunorubicin-based induction chemotherapy, and no patient received allogeneic hematopoietic stem cell transplantation (HSCT) during first complete remission (CR). For details regarding treatment protocols and sample collection, see Supplementary Information. All patients were enrolled on companion CALGB/Alliance protocols: 8461 (cytogenetic analyses), 9665 (tissue banking) and 20202 (molecular analyses).

Cytogenetic analyses were performed in institutional CALGB/Alliance cytogenetics laboratories. For the patient's karyotype to be considered normal, 20 metaphase cells from short-term cultures of pretreatment BM specimens had to have been analyzed and the normal result confirmed by central karyotype review.³³ All patients provided written informed consent for participation in these studies; study protocols were in accordance with the Declaration of Helsinki and approved by local Institutional Review Boards.

Single-gene expression analyses

The expression of *DNMT3B* transcript was assessed by NanoString nCounter assays (NanoString Technologies, Seattle, WA, USA; Supplementary Information).³⁴ These assays measured global expression of the *DNMT3B* gene, and did not allow for quantification of isoform-specific expression of *DNMT3B*. *DNMT3B* expression levels were normalized using *ABL* as an internal control. We also used NanoString nCounter assays to measure expression

of *BAALC*, *ERG* and *miR-155*, and real-time RT-PCR to measure *miR-3151* expression, all of which have been previously shown to affect prognosis of older CN-AML patients.^{10,15,16}

Mutational analyses

The presence or absence of *FLT3* internal tandem duplication (*FLT3*-ITD),^{35,36} *FLT3* tyrosine kinase domain mutations (*FLT3*-TKD),³⁷ *MLL* partial tandem duplication (*MLL*-PTD),³⁸ and mutations in the *NPM1*,⁵ *CEBPA*,³⁹ *WT1*,⁴⁰ *IDH1* and *IDH2*,⁷ *TET2*,⁴¹ *ASXL1*,⁸ *DNMT3A*⁹ and *RUNX1*⁴² genes were determined centrally as previously described.

Gene- and microRNA-expression profiling

The gene- and microRNA-expression profiling were assessed using the Affymetrix U133 plus 2.0 array (Affymetrix, Santa Clara, CA) and The Ohio State University (OSU) custom microRNA array (OSU_CCC Version 4.0), respectively, as previously reported,^{5,43} and detailed in the Supplementary Information. For *DNMT3B*, the Affymetrix U133 plus 2.0 arrays measured global *DNMT3B* expression levels, and did not quantify expression of the individual *DNMT3B* isoforms. For the gene- and microRNA-expression profiling, summary measures of gene and microRNA expression were computed, normalized, and filtered (Supplementary Information). A *DNMT3B* expression-associated signature (see Supplementary Information for details) was derived by comparing gene expression between high and low *DNMT3B* expressers in the Alliance cohort and in two additional sets of CN-AML patients with microarray and RNAseq gene expression data publicly available [German AML Cooperative Group (AMLCG)⁴⁴ and The Cancer Genome Atlas (TCGA)²⁵]. For comparison of the high *DNMT3B* expression signature with the *FLT3*-ITD signature we used gene set enrichment analysis (GSEA; for details see Supplementary Information). For the microRNA expression signature, only the CALGB/Alliance patients were used. Univariable significance levels of $P < 0.001$ (false discovery rates (FDR) < 0.01) were used to select genes and microRNAs that constituted the signatures. To assess enrichment of genes in the *DNMT3B* gene-expression-associated signature in distinct biologic processes, a Gene Ontology (GO) analysis was performed using the Database for Annotation, Visualization, and Integrated Discovery (DAVID).⁴⁵ We identified as statistically significant “annotation clusters” those clusters of GO terms with enrichment scores of > 2.0 , P -values < 0.001 and Benjamini corrected P -values < 0.05 . All molecular analyses were performed centrally at OSU.

DNA methylation

Genome-wide DNA methylation and levels of DNA methylation across the genome’s functional regions (i.e., genomic features) were measured using the MethylCap-seq assay as previously reported.¹⁸

Statistical analyses

The patients were dichotomized into high and low expressers using the median cut. This cut was supported by significant results of the trend test applied to outcome of patients divided into quartiles by *DNMT3B* expression ($P = 0.001$). We compared pretreatment features and outcome between patients with high and low *DNMT3B* expression. Definitions of clinical

endpoints [i.e., CR rates, disease-free (DFS) and overall (OS) survival] are provided in the Supplementary Information. Baseline characteristics between high and low *DNMT3B* expressers were compared using the Fisher's exact test for categorical and the Wilcoxon rank-sum test for continuous variables.⁴⁶ The categorical variables included the European LeukemiaNet (ELN) Genetic Groups.⁴⁷ The ELN guidelines classify CN-AML patients within the Favorable or Intermediate-I Genetic Groups based on *CEBPA*, *NPM1* and *FLT3* mutational status. The ELN Favorable Genetic Group consists of CN-AML patients with *CEBPA* mutation and/or *NPM1* mutation without *FLT3*-ITD, whereas the Intermediate-I Genetic Group is comprised of patients with wild-type *CEBPA* and *FLT3*-ITD with or without *NPM1* mutation, or wild-type *NPM1* without *FLT3*-ITD.⁴⁷

For time-to-event analyses, we calculated survival estimates using the Kaplan-Meier method, and compared groups by the log-rank test.⁴⁶ In order to provide the odds ratios and hazard ratios and associated confidence intervals, logistic regression and Cox proportional hazards models were generated to compare outcomes between high and low *DNMT3B* expressers for CR and survival endpoints (DFS, OS), respectively, and *P*-values from the Wald test are reported. We constructed multivariable logistic regression models to analyze factors associated with the achievement of CR, and multivariable Cox proportional hazards models for factors associated with survival endpoints,⁴⁶ the details of which are provided in the Supplementary Information. All analyses were performed by the Alliance for Clinical Trials in Oncology Statistics and Data Center.

RESULTS

Associations of DNMT3B expression with pretreatment clinical and molecular characteristics

At diagnosis, high *DNMT3B* expressers had higher white blood counts (WBC; $P=0.004$), and percentages of blood ($P=0.004$) and marrow ($P=0.02$) blasts than low *DNMT3B* expressers. Concerning molecular features, high *DNMT3B* expressers were more often *FLT3*-ITD-positive ($P<0.001$) and classified in the ELN Intermediate-I Genetic Group ($P=0.02$). *IDH2*-R140 mutations were less frequent in high *DNMT3B* expressers, whereas six of seven *IDH2*-R172 mutations were detected in this patient group. High *DNMT3B* expressers also had higher *ERG* ($P<0.001$), *BAALC* ($P=0.002$) and *miR-155* ($P=0.006$) expression than low expressers (Table 1, Supplementary Figure S1).

Associations of DNMT3B expression with clinical outcome in the entire patient cohort

With a median follow-up for patients alive of 5.1 years (range, 2.3–11.6 years), high *DNMT3B* expressers had lower CR rates ($P=0.002$, Wald test; 58% vs 78%), and shorter DFS ($P=0.02$, Wald test) and OS ($P<0.001$, Wald test) than *DNMT3B* low expressers (Table 2, Figure 1).

In a multivariable model for CR, *DNMT3B* expression remained prognostic ($P=0.04$), after adjustment for *BAALC* expression status ($P<0.001$), WBC ($P=0.007$) and age ($P=0.02$) (Table 3). High *DNMT3B* expressers were half as likely to achieve a CR as low expressers. In multivariable analysis for DFS, high *DNMT3B* expression associated with shorter DFS

($P=0.04$), once adjusted for *BAALC* expression ($P=0.004$), *DNMT3A*-R882 mutation status ($P=0.009$) and ELN Genetic Groups ($P=0.03$). The risk of experiencing relapse or death was 46% higher for high *DNMT3B* expressers than for low expressers. *DNMT3B* expression also remained prognostic for OS ($P=0.001$), after adjustment for *BAALC* expression ($P<0.001$), *miR-3151* ($P=0.02$) and *miR-155* ($P=0.02$). The risk of death was 72% higher for high *DNMT3B* expressers compared with low expressers (Table 3).

Associations of DNMT3B expression with clinical outcome in ELN genetic groups

We analyzed the associations of *DNMT3B* expression with outcome separately within the ELN Favorable and Intermediate-I Genetic Groups. Within the Favorable Group ($n=94$), there was no significant difference in CR rates (71% vs 84%, $P=0.14$, Wald test) or DFS ($P=0.10$, Wald test) between high and low *DNMT3B* expressers. However, high expressers had shorter OS ($P=0.002$, Wald test) than low expressers (Table 2, Figures 2A and 2B). In multivariable analyses for the ELN Favorable Genetic Group (Table 3), *DNMT3B* expression remained significant for OS ($P=0.003$) after adjustment for *BAALC* expression ($P=0.01$). High *DNMT3B* expressers were twice as likely to die as low expressers.

In the Intermediate-I Group ($n=108$), high *DNMT3B* expressers had a lower CR rate (49% vs 73%, $P=0.01$, Wald test) and shorter OS ($P=0.03$, Wald test) than low *DNMT3B* expressers, but there was no significant difference in DFS between the groups (Table 2, Figures 2C and 2D). In multivariable analyses within the ELN Intermediate-I Genetic Group (Table 3), *DNMT3B* expression was significant for OS ($P=0.02$), after adjustment for *BAALC* expression ($P=0.03$), *miR-3151* expression ($P=0.03$) and WBC ($P=0.03$). High *DNMT3B* expressers were 1.7 times more likely to die than low expressers.

Genome-wide gene-expression profiles associated with DNMT3B expression

To gain biologic insights into the role of *DNMT3B*, we derived a *DNMT3B*-associated gene-expression profile using three independent sets of CN-AML patients, i.e., CALGB/Alliance ($n=177$), AMLCG ($n=75$) and TCGA ($n=88$). We identified 195 upregulated genes and 168 downregulated genes that were significantly associated with higher *DNMT3B* expression in each of the three cohorts (Supplementary Table S1). Since high *DNMT3B* expression was associated with the presence of *FLT3*-ITD (Table 1), we performed GSEA to test whether a set of 195 genes that are upregulated in high *DNMT3B* expressers is associated with a set of genes differentially expressed between patients who harbored *FLT3*-ITD versus those who did not (Supplementary Information). We found a significant correlation between the high *DNMT3B* expression and *FLT3*-ITD signatures ($P=0.006$; FDR=0.006; Supplementary Figure S2). Among the genes upregulated in high *DNMT3B* expressers, we noted a variety of genes previously involved in AML including *CDK6* and *WT1* that encode cyclin kinase and transcription factor proteins, respectively. Among the downregulated genes, we noted genes involved with both normal monocyte/macrophage differentiation and immune function including *CD14*, *TLR4*, *CEBPB* and *TLR8*.

Gene Ontology was used to assess the biologic features of the *DNMT3B*-expression profile (Table 4). For *DNMT3B*-associated upregulated genes, there were three GO terms comprising genes involved in nucleotide biosynthetic processes and metabolism and

included in annotation cluster 1 that had a trend for statistical significance (Benjamini P -value <0.1). For *DNMT3B*-associated downregulated genes, cellular processes included lysosome biology, endocytosis and membrane signaling. These results may be interpreted as consistent with the previously noted dysregulated genes involved in monocyte/macrophage differentiation and activity.

Genome-wide microRNA profiles associated with DNMT3B expression

The influence of *DNMT3B* expression on microRNA genome-wide profiles could be evaluated in 162 patients. In contrast to coding genes, only four microRNAs were differentially expressed between high and low *DNMT3B* expressers ($P = 0.001$). High *DNMT3B* expression was associated with *miR-133b* upregulation, and *miR-148a*, *miR-122* and *miR-409-3p* downregulation. *miR-133b* upregulation in high *DNMT3B* expressers was somewhat surprising as this microRNA was reported to have tumor suppressor activity in other cancers.^{48,49} However, consistent with the downregulated gene-expression profile discussed above, *miR-133b* has recently been shown to target GM-CSF, a cytokine involved in granulocyte-monocyte/macrophage differentiation.⁵⁰ Among the downregulated microRNAs, *miR-148a* was reported to target *DNMT3B* and to be itself a target of aberrant hypermethylation in cancer.^{51,52} Lower expression of *miR-122* has been associated with aggressive hepatocellular carcinoma and *miR-409-3p* with cell invasion and metastasis in gastric cancer⁵³⁻⁵⁶; however, a role for these microRNAs in AML is currently unknown.

Genome-wide methylation profiling associated with DNMT3B expression

Since *DNMT3B* encodes a methyltransferase that mediates *de novo* DNA methylation, we assessed whether high and low *DNMT3B* expressers differed in DNA methylation patterns. Surprisingly, we found no significant differences in genome-wide DNA methylation levels or in the numbers of differentially methylated regions (DMRs)¹⁸ in distinct functional genomic regions (e.g., gene promoters) when high versus low *DNMT3B* expressers were compared.

DISCUSSION

In this study, we report that high *DNMT3B* expression associates with lower CR rates and shorter DFS and OS in chemotherapy-treated CN-AML patients aged ≥ 60 years. High *DNMT3B* expression was associated with such adverse prognostic factors as *FLT3*-ITD, high *ERG*, *BAALC* and *miR-155* expression and the ELN Intermediate-I Genetic Group; nevertheless the association of *DNMT3B* expression with clinical outcome is independent from the aforementioned and other established molecular and clinical prognosticators for all outcome endpoints studied.

Our findings are consistent to some extent with the only, to our knowledge, previous study that assessed the prognostic value of *DNMT3B* expression.²⁶ Although in the subset of 93 CN-AML patients, Hayette *et al.*²⁶ did not find significant differences in event-free survival (EFS) or OS between high and low *DNMT3B* expressers, high *DNMT3B* expressers had a shorter EFS than low *DNMT3B* expressers in the whole cytogenetically diverse cohort of 191 AML patients analyzed. It is difficult to directly compare their results with ours since

approximately one-half of the patients analyzed by Hayette *et al.*²⁶ had various abnormal karyotypes, more than two-thirds of the patients were younger than 60 years and a quarter underwent allogeneic HSCT in first CR. Thus, although the two studies are not comparable, they both conclude that higher *DNMT3B* expression is associated with worse outcome in AML.

Recently, the ELN Reporting System,⁴⁷ which for CN-AML is based on only three molecular markers (i.e., *FLT3-ITD*, *CEBPA* and *NPM1* mutations), was shown to provide important prognostic information in AML.⁵⁷ However, we and others have shown that additional molecular markers, such as *TET2*,⁴¹ *ASXL1*,⁸ *RUNX1*⁴² and *DNMT3A*⁵⁸ mutations and expression of *MNI*,¹² *miR-155*¹⁵ and *miR-3151*,¹⁶ may refine outcome prediction of CN-AML patients within the ELN Genetic Groups. Hence, in the current study, we investigated whether considering *DNMT3B* expression as a novel prognosticator could alter patient classification within the ELN Genetic Groups. In the Favorable Group, we found that low *DNMT3B* expression identified a subset of CN-AML patients with a significantly longer OS, thus making *DNMT3B* expression the third molecular marker, in addition to *ASXL1* mutations⁸ and *miR-155* expression,¹⁵ capable of refining prognostication of older patients in this ELN Genetic Group. We also observed a significant difference in OS between high and low *DNMT3B* expressers classified in the ELN Intermediate-I Genetic Group. Previously, *RUNX1* mutations⁴² and expression levels of *MNI*,¹² *miR-155*¹⁵ and *miR-3151*¹⁶ were demonstrated to add prognostic information in this ELN Genetic Group.

We report the first, to our knowledge, *DNMT3B*-associated gene- and microRNA-expression and DNA methylation profiles in CN-AML. We were able to derive a strong gene expression profile comprising 363 genes by overlapping the microarray results from three independent sets of patients. The profile was quite heterogeneous, comprising genes encoding for proteins involved in multiple biologic processes that play a role in leukemia cell differentiation, proliferation and survival. Among the downregulated genes, we noted enrichment of genes involved in monocyte/macrophage differentiation and activity, suggesting a role of *DNMT3B* in impairing differentiation of the leukemic blasts into cells with normal innate immunity activity. Using GSEA, we found a significant correlation between the high *DNMT3B* expression and *FLT3-ITD* signatures (Supplementary Figure S2). This, along with the increased frequency of *FLT3-ITD* in high *DNMT3B* expressers (Table 1), suggests the existence of a functional association between high expression of the *DNMT3B* gene and *FLT3-ITD*.

In contrast, the *DNMT3B*-associated microRNA profile was relatively weak, comprising only four microRNAs that were differentially expressed in high versus low *DNMT3B* expressers. Nevertheless, the unique upregulation of expression of *miR-133b*, recently reported to target GM-CSF,⁵⁰ in *DNMT3B* high expressers was somewhat consistent with the enrichment of the gene-expression profile in multiple downregulated genes involved in the differentiation and activity of hematopoietic cells participating in innate immunity.

Surprisingly, despite the fact that *DNMT3B* encodes a DNA methyltransferase, we observed no significant association of high *DNMT3B* levels and DNA methylation changes. No difference in global DNA methylation levels and number of DMRs could be identified

between *DNMT3B* high and low expressers. Our results are reminiscent of a recent report showing that changes in *DNMT3B* expression did not affect methylation levels of putative DNMT3B target genes.⁵⁹ Moreover, Russler-Germain *et al.*⁶⁰ have recently demonstrated that DNA methylation levels in leukemic blasts from CN-AML patients are not influenced by *DNMT3B* expression since mainly inactive splice variants of *DNMT3B* are expressed in these cells. Overall, therefore, these data may suggest that although overexpressed *DNMT3B* is a potentially valuable predictive marker for response to conventional chemotherapy, it does not necessarily identify subsets of older AML patients characterized by aberrant DNA methylation who might be responsive to hypomethylating azanucleosides.

In summary, we have demonstrated that *DNMT3B* expression constitutes an independent prognostic factor in older CN-AML patients treated intensively, and could also refine the ELN classification. Furthermore, we have provided some insights into the biologic activity of *DNMT3B* in CN-AML, which is seemingly independent from mechanisms of DNA hypermethylation and/or microRNA-dependent gene repression. Further studies focused on gaining more clinical and mechanistic insights into the leukemogenic role of *DNMT3B* expression are warranted to design active therapeutic strategies for high *DNMT3B* expressers in CN-AML.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The Cancer and Leukemia Group B institutions, and their principal investigators participating in this study are provided in the Supplementary Information. We thank Donna Bucci and the CALGB/Alliance Leukemia Tissue Bank at The Ohio State University Comprehensive Cancer Center, Columbus, OH, for sample processing and storage services; Lisa J. Sterling and Chris Finks for data management; and The Ohio State University Comprehensive Cancer Center's Nucleic Acid and Microarray Shared Resources for technical support. This work was supported in part by the National Cancer Institute (grants CA101140, CA114725, CA140158, CA31946, CA33601, CA16058, CA77658, and CA129657), the Coleman Leukemia Research Foundation, the Pelotonia Fellowship Program (A.-K.E.), the Conquer Cancer Foundation (J.H.M.) and the Deutsche Krebshilfe-Dr Mildred Scheel Cancer Foundation (H.B.).

REFERENCES

1. Mrózek K, Heerema NA, Bloomfield CD. Cytogenetics in acute leukemia. *Blood Rev.* 2004; 18:115–136. [PubMed: 15010150]
2. Mrózek K, Marcucci G, Paschka P, Whitman SP, Bloomfield CD. Clinical relevance of mutations and gene-expression changes in adult acute myeloid leukemia with normal cytogenetics: are we ready for a prognostically prioritized molecular classification? *Blood.* 2007; 109:431–448. [PubMed: 16960150]
3. Walker A, Marcucci G. Molecular prognostic factors in cytogenetically normal acute myeloid leukemia. *Expert Rev Hematol.* 2012; 5:547–558. [PubMed: 23146058]
4. Whitman SP, Maharry K, Radmacher MD, Becker H, Mrózek K, Margeson D, et al. FLT3 internal tandem duplication associates with adverse outcome and gene- and microRNA expression signatures in patients 60 years of age or older with primary cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *Blood.* 2010; 116:3622–3626. [PubMed: 20656931]
5. Becker H, Marcucci G, Maharry K, Radmacher MD, Mrózek K, Margeson D, et al. Favorable prognostic impact of *NPM1* mutations in older patients with cytogenetically normal de novo acute

- myeloid leukemia and associated gene- and microRNA-expression signatures: a Cancer and Leukemia Group B study. *J Clin Oncol.* 2010; 28:596–604. [PubMed: 20026798]
6. Becker H, Marcucci G, Maharry K, Radmacher MD, Mrózek K, Margeson D, et al. Mutations of the Wilms tumor 1 gene (*WT1*) in older patients with primary cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *Blood.* 2010; 116:788–792. [PubMed: 20442368]
 7. Marcucci G, Maharry K, Wu Y-Z, Radmacher MD, Mrózek K, Margeson D, et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol.* 2010; 28:2348–2355. [PubMed: 20368543]
 8. Metzeler KH, Becker H, Maharry K, Radmacher MD, Kohlschmidt J, Mrózek K, et al. *ASXL1* mutations identify a high-risk subgroup of older patients with primary cytogenetically normal AML within the ELN Favorable genetic category. *Blood.* 2011; 118:6920–6929. [PubMed: 22031865]
 9. Marcucci G, Metzeler KH, Schwind S, Becker H, Maharry K, Mrózek K, et al. Age-related prognostic impact of different types of *DNMT3A* mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J Clin Oncol.* 2012; 30:742–750. [PubMed: 22291079]
 10. Schwind S, Marcucci G, Maharry K, Radmacher MD, Mrózek K, Holland KB, et al. BAALC and ERG expression levels are associated with outcome and distinct gene- and microRNA expression profiles in older patients with de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *Blood.* 2010; 116:5660–5669. [PubMed: 20841507]
 11. Heuser M, Beutel G, Krauter J, Döhner K, von Neuhoff N, Schlegelberger B, et al. High meningioma 1 (MN1) expression as a predictor for poor outcome in acute myeloid leukemia with normal cytogenetics. *Blood.* 2006; 108:3898–3905. [PubMed: 16912223]
 12. Schwind S, Marcucci G, Kohlschmidt J, Radmacher MD, Mrózek K, Maharry K, et al. Low expression of MN1 associates with better treatment response in older patients with de novo cytogenetically normal acute myeloid leukemia. *Blood.* 2011; 118:4188–4198. [PubMed: 21828125]
 13. Marcucci G, Mrózek K, Radmacher MD, Garzon R, Bloomfield CD. The prognostic and functional role of microRNAs in acute myeloid leukemia. *Blood.* 2011; 117:1121–1129. [PubMed: 21045193]
 14. Schwind S, Maharry K, Radmacher MD, Mrózek K, Holland KB, Margeson D, et al. Prognostic significance of expression of a single microRNA, *miR-181a*, in cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol.* 2010; 28:5257–5264. [PubMed: 21079133]
 15. Marcucci G, Maharry KS, Metzeler KH, Volinia S, Wu Y-Z, Mrózek K, et al. Clinical role of microRNAs in cytogenetically normal acute myeloid leukemia: *miR-155* upregulation independently identifies high-risk patients. *J Clin Oncol.* 2013; 31:2086–2093. [PubMed: 23650424]
 16. Eisfeld AK, Marcucci G, Maharry K, Schwind S, Radmacher MD, Nicolet D, et al. *miR-3151* interplays with its host gene BAALC and independently affects outcome of patients with cytogenetically normal acute myeloid leukemia. *Blood.* 2012; 120:249–258. [PubMed: 22529287]
 17. Marcucci G, Yan P, Maharry K, Frankhouser D, Nicolet D, Metzeler KH, et al. Epigenetics meets genetics in acute myeloid leukemia: clinical impact of a novel seven-gene score. *J Clin Oncol.* 2014; 32:548–556. [PubMed: 24378410]
 18. Yan P, Frankhouser D, Murphy M, Tam HH, Rodriguez B, Curfman J, et al. Genome-wide methylation profiling in decitabine-treated patients with acute myeloid leukemia. *Blood.* 2012; 120:2466–2474. [PubMed: 22786882]
 19. Renneville A, Boissel N, Nibourel O, Berthon C, Helevaut N, Gardin C, et al. Prognostic significance of DNA methyltransferase 3A mutations in cytogenetically normal acute myeloid leukemia: a study by the Acute Leukemia French Association. *Leukemia.* 2012; 26:1247–1254. [PubMed: 22289988]
 20. Hervouet E, Vallette FM, Cartron PF. Dnmt3/transcription factor interactions as crucial players in targeted DNA methylation. *Epigenetics.* 2009; 4:487–499. [PubMed: 19786833]

21. Trowbridge JJ, Sinha AU, Zhu N, Li M, Armstrong SA, Orkin SH. Haploinsufficiency of *Dnmt1* impairs leukemia stem cell function through derepression of bivalent chromatin domains. *Genes Dev.* 2012; 26:344–349. [PubMed: 22345515]
22. Li KK, Luo LF, Shen Y, Xu J, Chen Z, Chen SJ. DNA methyltransferases in hematologic malignancies. *Semin Hematol.* 2013; 50:48–60. [PubMed: 23507483]
23. Wang J, Walsh G, Liu DD, Lee JJ, Mao L. Expression of *DNMT3B* variants and its association with promoter methylation of p16 and *RASSF1A* in primary non-small cell lung cancer. *Cancer Res.* 2006; 66:8361–8366. [PubMed: 16951144]
24. Hlady RA, Novakova S, Opavska J, Klinkebiel D, Peters SL, Bies J, et al. Loss of *Dnmt3b* function upregulates the tumor modifier *Ment* and accelerates mouse lymphomagenesis. *J Clin Invest.* 2012; 122:163–177. [PubMed: 22133874]
25. Cancer Genome Atlas Research Network. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *New Engl J Med.* 2013; 368:2059–2074. [PubMed: 23634996]
26. Hayette S, Thomas X, Jallades L, Chabane K, Charlot C, Tigaud I, et al. High DNA methyltransferase DNMT3B levels: a poor prognostic marker in acute myeloid leukemia. *PLoS One.* 2012; 7:e51527. [PubMed: 23251566]
27. Blum W, Garzon R, Klisovic RB, Schwind S, Walker A, Geyer S, et al. Clinical response and *miR-29b* predictive significance in older AML patients treated with a 10-day schedule of decitabine. *Proc Natl Acad Sci U S A.* 2010; 107:7473–7478. [PubMed: 20368434]
28. Mayer RJ, Davis RB, Schiffer CA, Berg DT, Powell BL, Schulman P, et al. Intensive postremission chemotherapy in adults with acute myeloid leukemia. *N Engl J Med.* 1994; 331:896–903. [PubMed: 8078551]
29. Stone RM, Berg DT, George SL, Dodge RK, Paciucci PA, Schulman P, et al. Granulocyte-macrophage colony-stimulating factor after initial chemotherapy for elderly patients with primary acute myelogenous leukemia. *N Engl J Med.* 1995; 332:1671–1677. [PubMed: 7760868]
30. Lee EJ, George SL, Caligiuri M, Sztatowski TP, Powell BL, Lemke S, et al. Parallel phase I studies of daunorubicin given with cytarabine and etoposide with or without the multidrug resistance modulator PSC-833 in previously untreated patients 60 years of age or older with acute myeloid leukemia: results of Cancer and Leukemia Group B study 9420. *J Clin Oncol.* 1999; 17:2831–2839. [PubMed: 10561359]
31. Baer MR, George SL, Sanford BL, Mrózek K, Kolitz JE, Moore JO, et al. Escalation of daunorubicin and addition of etoposide in the ADE regimen in acute myeloid leukemia patients aged 60 years and older: Cancer and Leukemia Group B study 9720. *Leukemia.* 2011; 25:800–807. [PubMed: 21321569]
32. Marcucci G, Moser B, Blum W, Stock W, Wetzler M, Kolitz JE, et al. A phase III randomized trial of intensive induction and consolidation chemotherapy ± oblimersen, a pro-apoptotic Bcl-2 antisense oligonucleotide in untreated acute myeloid leukemia patients >60 years old. *J Clin Oncol.* 2007; 25(suppl):360s. (abstract 7012).
33. Mrózek K, Carroll AJ, Maharry K, Rao KW, Patil SR, Pettenati MJ, et al. Central review of cytogenetics is necessary for cooperative group correlative and clinical studies of adult acute leukemia: the Cancer and Leukemia Group B experience. *Int J Oncol.* 2008; 33:239–244. [PubMed: 18636143]
34. Payton JE, Grieselhuber NR, Chang LW, Murakami M, Geiss GK, Link DC, et al. High throughput digital quantification of mRNA abundance in primary human acute myeloid leukemia samples. *J Clin Invest.* 2009; 119:1714–1726. [PubMed: 19451695]
35. Whitman SP, Archer KJ, Feng L, Baldus C, Becknell B, Carlson BD, et al. Absence of the wild-type allele predicts poor prognosis in adult de novo acute myeloid leukemia with normal cytogenetics and the internal tandem duplication of FLT3: a Cancer and Leukemia Group B study. *Cancer Res.* 2001; 61:7233–7239. [PubMed: 11585760]
36. Thiede C, Steudel C, Mohr B, Schaich M, Schäkel U, Platzbecker U, et al. Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. *Blood.* 2002; 99:4326–4335. [PubMed: 12036858]

37. Yamamoto Y, Kiyoi H, Nakano Y, Suzuki R, Kodera Y, Miyawaki S, et al. Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood*. 2001; 97:2434–2439. [PubMed: 11290608]
38. Whitman SP, Ruppert AS, Marcucci G, Mrózek K, Paschka P, Langer C, et al. Long-term disease-free survivors with cytogenetically normal acute myeloid leukemia and MLL partial tandem duplication: a Cancer and Leukemia Group B study. *Blood*. 2007; 109:5164–5167. [PubMed: 17341662]
39. Marcucci G, Maharry K, Radmacher MD, Mrózek K, Vukosavljevic T, Paschka P, et al. Prognostic significance of gene and microRNA expression signatures associated with, *CEBPA* mutations in cytogenetically normal acute myeloid leukemia with high-risk molecular features: a Cancer and Leukemia Group B study. *J Clin Oncol*. 2008; 26:5078–5087. [PubMed: 18809607]
40. Paschka P, Marcucci G, Ruppert AS, Whitman SP, Mrózek K, Maharry K, et al. Wilms' tumor 1 gene mutations independently predict poor outcome in adults with cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol*. 2008; 26:4595–4602. [PubMed: 18559874]
41. Metzeler KH, Maharry K, Radmacher MD, Mrózek K, Margeson D, Becker H, et al. TET2 mutations improve the new European LeukemiaNet risk classification of acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol*. 2011; 29:1373–1381. [PubMed: 21343549]
42. Mendler JH, Maharry K, Radmacher MD, Mrózek K, Becker H, Metzeler KH, et al. RUNX1 mutations are associated with poor outcome in younger and older patients with cytogenetically normal acute myeloid leukemia and with distinct gene and microRNA expression signatures. *J Clin Oncol*. 2012; 30:3109–3118. [PubMed: 22753902]
43. Marcucci G, Radmacher MD, Maharry K, Mrózek K, Ruppert AS, Paschka P, et al. MicroRNA expression in cytogenetically normal acute myeloid leukemia. *N Engl J Med*. 2008; 358:1919–1928. [PubMed: 18450603]
44. Metzeler KH, Hummel M, Bloomfield CD, Spiekermann K, Braess J, Sauerland MC, et al. An 86-probe-set gene-expression signature predicts survival in cytogenetically normal acute myeloid leukemia. *Blood*. 2008; 112:4193–4201. [PubMed: 18716133]
45. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*. 2009; 4:44–57. [PubMed: 19131956]
46. Vittinghoff, E.; Glidden, DV.; Shiboski, SC.; McCulloch, CE. *Regression Methods in Biostatistics: Linear, Logistic, Survival and Repeated Measures Models*. New York, NY, USA: Springer; 2005.
47. Döhner H, Estey EH, Amadori S, Appelbaum FR, Büchner T, Burnett AK, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010; 115:453–474. [PubMed: 19880497]
48. Zhao H, Li M, Li L, Yang X, Lan G, Zhang Y. MiR-133b is down-regulated in human osteosarcoma and inhibits osteosarcoma cells proliferation, migration and invasion, and promotes apoptosis. *PLoS One*. 2013; 8:e83571. [PubMed: 24391788]
49. Duan F-T, Qian F, Fang K, Lin K-Y, Wang W-T, Chen Y-Q. miR-133b, a muscle-specific microRNA, is a novel prognostic marker that participates in the progression of human colorectal cancer via regulation of CXCR4 expression. *Mol Cancer*. 2013; 12:164. [PubMed: 24330809]
50. Sturrock A, Mir-Kasimov M, Baker J, Rowley J, Paine R 3rd. Key role of microRNA in the regulation of granulocyte macrophage colony-stimulating factor expression in murine alveolar epithelial cells during oxidative stress. *J Biol Chem*. 2014; 289:4095–4105. [PubMed: 24371146]
51. Duursma AM, Kedde M, Schrier M, le Sage C, Agami R. miR-148 targets human DNMT3b protein coding region. *RNA*. 2008; 14:872–877. [PubMed: 18367714]
52. Zhu A, Xia J, Zuo J, Jin S, Zhou H, Yao L, et al. MicroRNA-148a is silenced by hypermethylation and interacts with DNA methyltransferase 1 in gastric cancer. *Med Oncol*. 2012; 29:2701–2709. [PubMed: 22167392]
53. Hsu SH, Wang B, Kota J, Yu J, Costinean S, Kutay H, et al. Essential metabolic, anti-inflammatory, and anti-tumorigenic functions of miR-122 in liver. *J Clin Invest*. 2012; 122:2871–2883. [PubMed: 22820288]

54. Köberle V, Kronenberger B, Pleli T, Trojan J, Imelmann E, Peveling-Oberhag J, et al. Serum microRNA-1 and microRNA-122 are prognostic markers in patients with hepatocellular carcinoma. *Eur J Cancer*. 2013; 49:3442–3449. [PubMed: 23810247]
55. Li A, Song W, Qian J, Li Y, He J, Zhang Q, et al. MiR-122 modulates type I interferon expression through blocking suppressor of cytokine signaling 1. *Int J Biochem Cell Biol*. 2013; 45:858–865. [PubMed: 23348614]
56. Zheng B, Liang L, Huang S, Zha R, Liu L, Jia D, et al. MicroRNA-409 suppresses tumour cell invasion and metastasis by directly targeting radixin in gastric cancers. *Oncogene*. 2012; 31:4509–4516. [PubMed: 22179828]
57. Mrózek K, Marcucci G, Nicolet D, Maharry KS, Becker H, Whitman SP, et al. Prognostic significance of the European LeukemiaNet standardized system for reporting cytogenetic and molecular alterations in adults with acute myeloid leukemia. *J Clin Oncol*. 2012; 30:4515–4523. [PubMed: 22987078]
58. Gaidzik VI, Schlenk RF, Paschka P, Stölzle A, Späth D, Kuendgen A, et al. Clinical impact of DNMT3A mutations in younger adult patients with acute myeloid leukemia: results of the AML Study Group (AML5SG). *Blood*. 2013; 121:4769–4777. [PubMed: 23632886]
59. Hagemann S, Kuck D, Stresemann C, Prinz F, Brueckner B, Mund C, et al. Antiproliferative effects of DNA methyltransferase 3B depletion are not associated with DNA demethylation. *PLoS One*. 2012; 7:e36125. [PubMed: 22563479]
60. Russler-Germain DA, Spencer DH, Young MA, Lamprecht TL, Miller CA, Fulton R, et al. The R882H DNMT3A mutation associated with AML dominantly inhibits wild-type DNMT3A by blocking its ability to form active tetramers. *Cancer Cell*. 2014; 25:442–454. [PubMed: 24656771]






Figure 1. Clinical outcome of CN-AML patients with high and low *DNMT3B* expression. Kaplan-Meier survival curves for (a) disease-free survival and (b) overall survival. *P*-values presented are from the log rank test.

a



b






Figure 2. Clinical outcome of CN-AML patients with high and low *DNMT3B* expression classified into European LeukemiaNet (ELN) Genetic Groups. Kaplan-Meier survival curves for (a) disease-free survival and (b) overall survival of patients in the ELN modified Favorable Genetic Group; (c) disease-free survival and (d) overall survival of patients in the ELN Intermediate-I Genetic Group. *P*-values presented are from the log rank test.

Table 1

Comparison of clinical and molecular characteristics of patients with cytogenetically normal acute myeloid leukemia with high versus low *DNMT3B* expression

Characteristic	High DNMT3B (n=105)	Low DNMT3B (n=105)	P
<i>Age, years</i>			0.82
Median	68	68	
Range	60–83	60–81	
<i>Sex, n (%)</i>			0.58
Male	58 (55)	53 (50)	
Female	47 (45)	52 (50)	
<i>Race, n (%)</i>			0.48
White	96 (92)	92 (89)	
Nonwhite	8 (8)	11 (11)	
<i>Hemoglobin, g/dl</i>			0.71
Median	9.4	9.3	
Range	6.5–12.4	5.4–15.0	
<i>Platelet count, × 10⁹/l</i>			0.87
Median	68	71	
Range	4–850	11–510	
<i>WBC, × 10⁹/l</i>			0.004
Median	43.7	21.8	
Range	1.0–450.0	0.8–249.3	
<i>Blood blasts, %</i>			0.004
Median	64	40	
Range	0–99	0–97	
<i>Bone marrow blasts, %</i>			0.02
Median	72	64	
Range	21–97	4–97	
<i>Extramedullary involvement, n (%)</i>	27 (27)	24 (23)	0.63
<i>NPM1, n (%)</i>			0.31
Mutated	67 (66)	60 (59)	
Wild-type	34 (34)	42 (41)	
<i>FLT3-ITD, n (%)</i>			<0.001
Present	54 (53)	21 (21)	
Absent	48 (47)	81 (79)	
<i>CEBPA, n (%)</i>			0.83 ^a

Characteristic	High DNMT3B (n=105)	Low DNMT3B (n=105)	P
Mutated	13 (13)	12 (12)	
Single mutated	10	5	
Double mutated	3	7	
Wild-type	88 (87)	90 (88)	
<i>ELN Genetic Group^b, n (%)</i>			0.02
Modified Favorable	38 (38)	56 (55)	
Intermediate-I	63 (62)	45 (45)	
<i>FLT3-TKD, n (%)</i>			0.83
Present	12 (12)	11 (11)	
Absent	89 (88)	91 (89)	
<i>WT1, n (%)</i>			0.41
Mutated	8 (8)	5 (5)	
Wild-type	93 (92)	97 (95)	
<i>TET2, n (%)</i>			1.00
Mutated	32 (32)	31 (32)	
Wild-type	68 (68)	67 (68)	
<i>MLL-PTD, n (%)</i>			1.00
Present	5 (6)	5 (6)	
Absent	74 (94)	81 (94)	
<i>IDH1, n (%)</i>			0.18
Mutated	14 (14)	8 (8)	
Wild-type	86 (86)	94 (92)	
<i>IDH2, n (%)</i>			0.05
Mutated	18 (18)	31 (30)	
R140	12	30	0.005 ^c
R172	6	1	0.13 ^d
Wild-type	82 (82)	71 (70)	
<i>RUNX1, n (%)</i>			0.31
Mutated	17 (18)	11 (12)	
Wild-type	79 (82)	81 (88)	
<i>ASXL1, n (%)</i>			0.69
Mutated	13 (13)	15 (15)	
Wild-type	87 (87)	84 (85)	
<i>DNMT3A</i>			0.65
Mutated	35 (35)	31 (32)	
R882	19	20	1.00 ^e

Characteristic	High DNMT3B (n=105)	Low DNMT3B (n=105)	P
Non-R882	16	11	0.40 ^f
Wild-type	64 (65)	66 (68)	
<i>ERG</i> expression group, ^{g,h} n (%)			<0.001
High	65 (62)	40 (38)	
Low	40 (38)	65 (62)	
<i>BAALC</i> expression group, ^{g,h} n (%)			0.002
High	64 (61)	41 (39)	
Low	41 (39)	64 (61)	
<i>miR-155</i> expression group, ^{g,h} n (%)			0.006
High	63 (60)	42 (40)	
Low	42 (40)	63 (60)	
<i>miR-3151</i> expression group, ^{g,i} n (%)			0.76
High	42 (49)	39 (46)	
Low	43 (51)	46 (54)	

Abbreviations: ELN, European LeukemiaNet; *FLT3*-ITD, internal tandem duplication of the *FLT3* gene; *FLT3*-TKD, tyrosine kinase domain mutation in the *FLT3* gene; *MLL*-PTD, partial tandem duplication of the *MLL* gene; n, number; WBC, white blood count.

^aThe *P*-value pertains to a comparison of frequencies of *CEBPA* mutations (single and double combined) versus *CEBPA* wild-type between high and low *DNMT3B* expressers.

^bThe ELN modified Favorable Genetic Group is defined as CN-AML patients with mutated *CEBPA* and/or mutated *NPM1* without *FLT3*-ITD. All remaining CN-AML patients (i.e., those with wild-type *CEBPA* and wild-type *NPM1* with or without *FLT3*-ITD, or mutated *NPM1* with *FLT3*-ITD) belong to the ELN Intermediate-I Genetic Group.⁴⁷

^cThe *P*-value pertains to a comparison of frequencies of *IDH2*-R140 mutations versus *IDH2* wild-type between high and low *DNMT3B* expressers.

^dThe *P*-value pertains to a comparison of frequencies of *IDH2*-R172 mutations versus *IDH2* wild-type between high and low *DNMT3B* expressers.

^eThe *P*-value pertains to a comparison of frequencies of *DNMT3A*-R882 mutations versus *DNMT3A* wild-type between high and low *DNMT3B* expressers.

^fThe *P*-value pertains to a comparison of frequencies of *DNMT3A* non-R882 mutations versus *DNMT3A* wild-type between high and low *DNMT3B* expressers.

^gThe median expression value was used as a cut point.

^hData was assessed by the NanoString nCounter assay.

ⁱData was assessed by real-time RT-PCR.

Table 2

Outcomes of patients with cytogenetically normal acute myeloid leukemia according to *DNMT3B* expression status

Endpoint	High DNMT3B (I)	Low DNMT3B (II)	<i>p</i> ^a	OR/HR (95% CI) I vs II
All patients	n=105	n=105		
<i>Complete remission, n (%)</i>	61 (58)	82 (78)	0.002	0.39 (0.21–0.71)
<i>Disease-free survival</i>			0.02	1.55 (1.09–2.20)
Median, years	0.6	1.1		
% Disease-free at 3 years (95% CI)	13 (6–23)	22 (14–31)		
% Disease-free at 5 years (95% CI)	11 (5–21)	15 (8–23)		
<i>Overall survival</i>			<0.001	1.85 (1.38–2.47)
Median, years	0.8	1.5		
% Alive at 3 years (95% CI)	11 (6–18)	29 (20–37)		
% Alive at 5 years (95% CI)	8 (4–14)	21 (14–29)		
Patients in the ELN modified Favorable Genetic Group ^b	n=38	n=56		
<i>Complete remission, n (%)</i>	27 (71)	47 (84)	0.14	0.47 (0.17–1.28)
<i>Disease-free survival</i>			0.10	1.54 (0.92–2.57)
Median, years	0.9	1.3		
% Disease-free at 3 years (95% CI)	15 (5–30)	30 (18–43)		
% Disease-free at 5 years (95% CI)	11 (3–26)	21 (11–34)		
<i>Overall survival</i>			0.002	2.04 (1.29–3.22)
Median, years	1.3	2.1		
% Alive at 3 years (95% CI)	18 (7–31)	39 (27–52)		
% Alive at 5 years (95% CI)	12 (4–24)	30 (19–43)		
Patients in the ELN Intermediate-1 Genetic Group ^b	n=63	n=45		
<i>Complete remission, n(%)</i>	31 (49)	33 (73)	0.01	0.35 (0.15–0.80)
<i>Disease-free survival</i>			0.25	1.36 (0.81–2.27)
Median, years	0.5	0.7		
% Disease-free at 3 years (95% CI)	13 (4–27)	9 (2–22)		
% Disease-free at 5 years (95% CI)	13 (4–27)	3 (0–13)		
<i>Overall survival</i>			0.03	1.57 (1.06–2.33)
Median, years	0.6	1.1		
% Alive at 3 years (95% CI)	8 (3–16)	16 (7–28)		
% Alive at 5 years (95% CI)	6 (2–14)	9 (3–19)		

Abbreviations: CI, confidence interval; ELN, European Leukemia Net; HR, hazard ratio; n, number; OR, odds ratio.

^a *P*-values provided are generated by logistic regression and Cox proportional hazards models to compare outcome of patients for CR and survival endpoints (DFS, OS), respectively, using the Wald test.

^b The ELN modified Favorable Genetic Group is defined as CN-AML patients with mutated *CEBPA* and/or mutated *NPM1* without *FLT3*-ITD. All remaining CN-AML patients (i.e., those with wild-type *CEBPA* and wild-type *NPM1* with or without *FLT3*-ITD, or mutated *NPM1* with *FLT3*-ITD) belong to the ELN Intermediate-I Genetic Group.⁴⁷

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Multivariable analyses of CN-AML patients according to DNMT3B expression status

Variable	Complete remission			Disease-free survival			Overall survival		
	OR (95% CI)	P	HR (95% CI)	HR (95% CI)	P	HR (95% CI)	HR (95% CI)	P	
All patients									
DNMT3B expression, high vs low	0.49 (0.25–0.97)	0.04	1.47 (1.01–2.13)	0.04	1.72 (1.24–2.38)	0.001			
BAALC expression, high vs low	0.21 (0.10–0.43)	<0.001	1.79 (1.21–2.65)	0.004	1.98 (1.39–2.80)	<0.001			
WBC, per 50-unit increase	0.69 (0.52–0.90)	0.007							
Age, per 10-year increase	0.49 (0.27–0.89)	0.02							
DNMT3A ^a									
R882 mutated vs wild-type			1.85 (1.17–2.95)	0.009					
non-R882 mutated vs wild-type			0.92 (0.52–1.61)	0.76					
ELN Genetic Group, Favorable vs Intermediate-1 ^b			0.65 (0.45–0.96)	0.03					
miR-3151 expression, high vs low						1.51 (1.07–2.13)	0.02		
miR-155 expression, high vs low						1.47 (1.06–2.05)	0.02		
Patients in the ELN modified Favorable Genetic Group ^b									
DNMT3B expression, high vs low						1.99 (1.26–3.15)	0.003		
BAALC expression, high vs low						1.81 (1.13–2.92)	0.01		
Patients in the ELN Intermediate-1 Genetic Group ^b									
DNMT3B expression, high vs low						1.73 (1.10–2.72)	0.02		
BAALC expression, high vs low						1.88 (1.05–3.35)	0.03		
miR-3151 expression, high vs low						1.79 (1.05–3.07)	0.03		

No models including a significant term for DNMT3B expression were found

No models including a significant term for DNMT3B expression were found

Variable	Complete remission		Disease-free survival		Overall survival	
	OR (95% CI)	P	HR (95% CI)	P	HR (95% CI)	P
WBC, per 50-unit increase					1.16 (1.01–1.33)	0.03

Abbreviations: CI, confidence interval; ELN, European LeukemiaNet; HR, hazard ratio; OR, odds ratio; WBC, white blood count. An odds ratio less than 1 means a lower CR rate for the higher values of the continuous variables and the first category listed for the categorical variables. A hazard ratio greater than 1 (less than 1) corresponds to a higher (lower) risk of an event for higher values of continuous variables and the first category listed of a dichotomous variable. Variables were considered for inclusion in the multivariable models if they had a univariable *P*-value of 0.20. See the Supplementary Information for a full list of variables evaluated in univariable analyses. Since *NPM1*, *FLT3-ITD*, and *CEBPA* mutations are integrated in the ELN genetic classification, they were not additionally considered as individual variables. In the entire patient cohort, variables considered for inclusion in the model for achievement of CR were *DNMT3B*, *ERG*, *BAALC*, *miR-155* and *miR-3151* expression, ELN Genetic Groups, *WT1* and *ASXL1* mutation status, WBC, age and extramedullary involvement. In the model for DFS, we considered *DNMT3B*, *ERG*, *BAALC* and *miR-3151* expression, ELN Genetic Groups, *FLT3-TKD*, *ASXL1*, *DNMT3A* non-R882 mutation status and extramedullary involvement; and in the model for OS, *DNMT3B*, *ERG*, *BAALC*, *miR-155* and *miR-3151* expression, ELN Genetic Groups, *MLL-PTD*, *WT1*, *ASXL1*, *DNMT3A* non-R882 mutation status, WBC and extramedullary involvement. For patients in the modified Favorable ELN Genetic Group, variables considered for inclusion in the model for OS were *DNMT3B*, *ERG*, *BAALC* and *miR-155* expression, *ASXL1* and *TET2* mutation status and extramedullary involvement. For patients in the Intermediate-I ELN Genetic Group, variables considered for inclusion in the model for OS were *DNMT3B*, *ERG*, *BAALC*, *miR-155* and *miR-3151* expression, *RUNX1*, *IDH1*, *DNMT3A* non-R882 mutation status, and WBC and hemoglobin.

^aThe types of *DNMT3A* mutations detected in our cohort are provided in Supplementary Table S2.

^bThe ELN modified Favorable Genetic Group is defined as CN-AML patients with mutated *CEBPA* and/or mutated *NPM1* without *FLT3-ITD*. All remaining CN-AML patients (i.e., those with wild-type *CEBPA* and wild-type *NPM1* with or without *FLT3-ITD*, or mutated *NPM1* with *FLT3-ITD*) belong to the ELN Intermediate-I Genetic Group.⁴⁷

Table 4Gene Ontology terms associated with differentially expressed genes in the high *DNMT3B* expression group

Biologic process or cellular component	Number of genes	Benjamini P-value
Associations with genes upregulated in the high <i>DNMT3B</i> expression group		
Annotation Cluster 1.		
GOTERM_BP: nucleotide biosynthetic process	10	0.098
GOTERM_BP: nucleobase, nucleoside and nucleotide biosynthetic process	10	0.066
GOTERM_BP: nucleobase, nucleoside, nucleotide and nucleic acid biosynthetic process	10	0.066
Associations with genes downregulated in the high <i>DNMT3B</i> expression group		
Annotation Cluster 1		
GOTERM_CC: vacuole	12	0.0052
GOTERM_CC: lytic vacuole	10	0.012
GOTERM_CC: lysosome	10	0.012
Annotation Cluster 2		
GOTERM_BP: endocytosis	11	0.0091
GOTERM_BP: membrane invagination	11	0.0091
GOTERM_BP: membrane organization	12	0.089
Annotation Cluster 3		
GOTERM_CC: intrinsic to membrane	74	0.028

Abbreviations: GO, Gene Ontology, see also Huang da *et al.*⁴⁵; BP, biologic process; CC, cellular component.