

2014

A rare case of double-chambered right ventricle associated with ventricular septal defect and congenital absence of the pulmonary valve

G. Khoueiry
Northwell Health

T. Bhat
Northwell Health

M. Tantray
Northwell Health

M. Meghani
Northwell Health

N. Abi Rafeh
Northwell Health

See next page for additional authors

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>

 Part of the [Cardiology Commons](#)

Recommended Citation

Khoueiry G, Bhat T, Tantray M, Meghani M, Abi Rafeh N, Abdallah M, Hoyek W. A rare case of double-chambered right ventricle associated with ventricular septal defect and congenital absence of the pulmonary valve. . 2014 Jan 01; 8():Article 2436 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/2436>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

Authors

G. Khoueiry, T. Bhat, M. Tantray, M. Meghani, N. Abi Rafeh, M. Abdallah, and W. Hoyek

SHORT REPORT

OPEN ACCESS

Full open access to this and
thousands of other papers at
<http://www.la-press.com>.

A Rare Case of Double-Chambered Right Ventricle Associated with Ventricular Septal Defect and Congenital Absence of the Pulmonary Valve

Georges Khoueiry¹, Tariq Bhat¹, Mohmad Tantray², Mustafain Meghani², Nidal Abi Rafeh¹, Mokhtar Abdallah² and Wissam Hoyek¹

¹Department of Cardiology, Staten Island University Hospital, Staten Island, NY. ²Department of Medicine, Staten Island University Hospital, Staten Island, NY. Corresponding author email: mohivddin_bhat@yahoo.com

Abstract: Double-chambered right ventricle (DCRV) is a rare congenital heart disorder involving 2 different right ventricle (RV) pressure compartments that is often associated with ventricular septal defect (VSD). Usually, the obstruction is caused by an anomalous muscle bundle crossing the RV from the interventricular septum to the RV free wall. We are reporting a case of double-chambered right ventricle associated with ventricular septal defect and congenital absence of the pulmonary valve, a rare form of congenital infundibular pulmonary stenosis. In addition to ventricular septal defect, our patient had congenital absence of the pulmonary valve, which is very unusual and has never been reported to our knowledge.

Keywords: cardiology, magnetic resonance imaging, echocardiography, doubled-chambered right ventricle, ventricular septal defect

Clinical Medicine Insights: Circulatory, Respiratory and Pulmonary Medicine 2014:8 1–3

doi: [10.4137/CCRPM.S11174](https://doi.org/10.4137/CCRPM.S11174)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article published under the Creative Commons CC-BY-NC 3.0 license.

Case Presentation

A 45-year-old female presented to the cardiology office complaining of persistent dyspnea on exertion. She also reported intermittent chest pain along with bilateral lower extremity edema. Symptoms had been present for past 2 years and were becoming progressively worse. Her past medical history was consistent with hypothyroidism. There was no evidence of deep vein thrombosis on lower extremity duplex. A transthoracic echocardiogram showed a normal left ventricular function and a small ventricular septal defect (VSD) (Video 1). Color Doppler analysis using transesophageal echocardiogram revealed turbulence in the proximal portion of right ventricle with a high velocity jet (Image 1A, Video 2). An abnormal muscle bundle was also evident separating the proximal from the distal infundibular chamber of right ventricle (Image 1B and C, Video 2). Measurement of the pressure gradient between the proximal and distal chamber was not done since the ultrasound beam could not be aligned with the jet. Magnetic resonance imaging (MRI) confirmed the finding of membranous ventricular septal defect without significant shunting and increased trabecular markings of the

subfundibular region of the right ventricle consistent with double-chambered right ventricle (DCRV) (Image 1D). The pulmonary valve was not seen on echocardiogram or MRI.

Discussion

DCRV is a rare form of congenital infundibular pulmonary stenosis. It is characterized by aberrant hypertrophied muscle bands that divide the right ventricle into 2 cavities with a pressure gradient across. Several subtypes of divided RV have been described.¹ Associated defects are present in approximately 80% to 90% of patients. VSD that involves the membranous septum is the most common associated congenital abnormality. Other frequent associated lesions include pulmonary valve stenosis and discrete subaortic stenosis. Vogel et al described 36 patients with membranous VSD and double-chambered right ventricle, 88% of who had echocardiographic evidence of subaortic stenosis, with evidence of progressive left ventricular outflow tract obstruction.²

Congenital absence of the pulmonary valve is uncommon and usually associated with VSD and




Figure 1. (A) A transesophageal echocardiogram with color doppler (short axis midesophageal view) showing turbulence in the proximal portion of right ventricle with a high velocity jet. (B) In the same view, the muscle bands are clearly illustrated (white arrow) and appear to divide the right ventricle into a proximal and distal infundibular chamber. (C) A modified transesophageal view also showing the muscle bands (white arrow). (D) MRI image (horizontal long axis view) showing increased trabecular markings and anomalous muscle bundles (arrow) dividing the right ventricle into a proximal and a distal chamber.



obstructive subvalvular pulmonary ring but not typically with DCRV.

Author Contributions

Analyzed the data: GK. Wrote the first draft of the manuscript: MM, MA. Contributed to the writing of the manuscript: TB, MT. Agree with manuscript results and conclusions: WH. Jointly developed the structure and arguments for the paper: TB, GK, NAR. Made critical revisions and approved final version: TB. All authors reviewed and approved of the final manuscript.

Funding

Author(s) disclose no funding sources.

Competing Interests

Author(s) disclose no potential conflicts of interest.

Disclosures and Ethics

As a requirement of publication the authors have provided signed confirmation of their compliance with

ethical and legal obligations including but not limited to compliance with ICMJE authorship and competing interests guidelines, that the article is neither under consideration for publication nor published elsewhere, of their compliance with legal and ethical guidelines concerning human and animal research participants (if applicable), and that permission has been obtained for reproduction of any copyrighted material. This article was subject to blind, independent, expert peer review. The reviewers reported no competing interests.

References

1. Restivo A, Cameron AH, Anderson RH, Allwork SP. Divided right ventricle: a review of its anatomical varieties. *Pediatr Cardiol.* 1984;5(3):197–204.
2. Vogel M, Smallhorn JF, Freedom RM, Coles J, Williams WG, Trusler GA. An echocardiographic study of the association of ventricular septal defect and right ventricular muscle bundles with a fixed subaortic abnormality. *Am J Cardiol.* 1988;61(10):857–60.