

2014

Genotyping of immune-related genetic variants identifies TYK2 as a novel associated locus for idiopathic inflammatory myopathies

M. Jani

J. Massey

L. R. Wedderburn

J. Vencovsky

K. Danko

See next page for additional authors

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>

 Part of the [Medical Molecular Biology Commons](#)

Recommended Citation

Jani M, Massey J, Wedderburn L, Vencovsky J, Danko K, Lundberg I, Padyukov L, Lee A, Gregersen P, Eumyonet , . Genotyping of immune-related genetic variants identifies TYK2 as a novel associated locus for idiopathic inflammatory myopathies. . 2014 Jan 01; 73(9):Article 2887 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/2887>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

Authors

M. Jani, J. Massey, L. R. Wedderburn, J. Vencovsky, K. Danko, I. E. Lundberg, L. Padyukov, A. Lee, P. K. Gregersen, Eumyonet, and +10 additional authors

Published in final edited form as:

Ann Rheum Dis. 2014 September ; 73(9): 1750–1752. doi:10.1136/annrheumdis-2014-205440.

Genotyping of immune-related genetic variants identifies *TYK2* as a novel associated locus for idiopathic inflammatory myopathies

M Jani¹, J Massey^{1,2}, L R Wedderburn³, J Vencovsky⁴, K Danko⁵, I E Lundberg⁶, L Padyukov⁶, A Selva-O'Callaghan⁷, T Radstake⁸, H Platt⁹, R B Warren², C E Griffiths², A Lee¹⁰, P K Gregersen¹⁰, F W Miller¹¹, W E Ollier⁹, R G Cooper¹², H Chinoy^{1,9}, J A Lamb⁹, and EUMYONET

¹Arthritis Research UK Centre for Epidemiology, Musculoskeletal Research Group, Manchester Academic Health Science Centre, University of Manchester, Manchester, UK ²The Dermatology Centre, Salford Royal NHS Foundation Trust, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK ³Rheumatology Unit, Institute of Child Health, University College London, London, UK ⁴Institute of Rheumatology, Charles University in Prague, Prague, Czech Republic ⁵University of Debrecen, Debrecen, Hungary ⁶Rheumatology Unit, Department of Medicine, Karolinska University Hospital, Solna, Karolinska Institutet, Stockholm, Sweden ⁷Vall d'Hebron General Hospital, Barcelona, Spain ⁸Laboratory of Translational Immunology, University Medical Center Utrecht, Utrecht, The Netherlands ⁹Centre for Integrated Genomic Medical Research, Manchester Academic Health Science Centre, University of Manchester, Manchester, UK ¹⁰Feinstein Institute for Medical Research, Manhasset, New York, USA ¹¹National Institute of Environmental Health Sciences, National Institutes of Health, Bethesda, Maryland, USA ¹²Department of Musculoskeletal Biology, University of Liverpool, Liverpool, UK

Idiopathic inflammatory myopathies (IIMs) may present as a primary autoimmune disorder, or overlap with other autoimmune/connective tissue diseases. The aetiology of IIM likely includes interactions between genetic and environmental factors. Several genetic variants common to multiple autoimmune disorders have been identified in recent genome-wide association studies (GWAS). A Myositis Genetics Consortium dermatomyositis (DM) GWAS also suggests genetic overlap with other autoimmune disorders.¹ We sought to

Copyright Article author (or their employer) 2014.

Produced by BMJ Publishing Group Ltd (& EULAR) under licence. To request permissions go to: <http://group.bmj.com/group/rights-licensing/permissions> To order reprints go to: <http://journals.bmj.com/cgi/reprintform>

Correspondence to: Dr Meghna Jani, Arthritis Research UK Centre for Epidemiology, Centre for Musculoskeletal Research, Institute of Inflammation and Repair, The University of Manchester, Manchester Academic Health Science Centre, Stopford Building, Oxford Road, Manchester M13 9PT, UK; Meghna.jani@manchester.ac.uk
HC and JAL are joint last authors.

Contributors MJ, HC, RGC and JAL were involved in the conception of the project. MJ, JM and JAL performed the methods and analysis of the project. The rest of the coauthors contributed samples for the purposes of the study, were involved in the discussion of final content and review/editing of the manuscript before submission.

Competing interests None.

Ethics approval Local ethics committees for each city.

Provenance and peer review Not commissioned; externally peer reviewed.

extend these findings to identify novel genetic risk factors in a large cohort of adult/juvenile patients with DM and polymyositis (PM), by genotyping immune-related single nucleotide polymorphisms (SNPs) not captured through the DM GWAS.¹

SNPs significantly associated ($p < 5 \times 10^{-8}$) with 10 autoimmune disorders (systemic lupus erythematosus, rheumatoid arthritis, juvenile idiopathic arthritis, coeliac disease, Crohn's disease, ulcerative colitis, psoriasis, type 1 diabetes, multiple sclerosis and systemic sclerosis) were identified from published GWAS or the National Human Genome Research Institute GWAS catalogue.² Unique SNPs were identified ($n=233$), of which 99 had not been directly genotyped or captured ($r^2 < 0.8$ with genotyped SNPs) through our DM GWAS.¹ These 99 SNPs were genotyped using Sequenom in 1001 European Caucasian individuals with definite/probable adult or juvenile DM or PM.³⁴ Eighty-three SNPs passed quality control. GWAS data² were imputed to the 1000G_phase 1 integrated_v3 reference panel using IMPUTE2.⁵ Concordance rates were $>98\%$ for 444/1001 cases overlapping the GWAS data set. INFO scores for the 83 SNPs were >0.6 . The 1001 Sequenom cases were merged with 168 unique cases from the GWAS. Samples with $>5\%$ missing data were excluded, resulting in 1149 cases and 3572 controls (table 1). Association tests were performed for the 83 SNPs in DM and PM subgroups including juvenile cases, and the overall IIM group (SNPTEST V.2.4.1). Imputed genotype uncertainty was accounted for using the 'score' method. A random-effects meta-analysis of the individual country data sets was performed (META V.1.5). Local research ethics committee approval and informed consent were obtained.

Outside the major histocompatibility complex (MHC), a non-synonymous SNP rs2304256 in *TYK2* was identified reaching Bonferroni corrected significance in DM and the overall IIM group, but not in the PM subgroup ($p=0.17$) (table 2). *TYK2* encodes a member of the Janus tyrosine kinase protein family, contributing to cytokine receptor signaling via its catalytic activity or kinase-independent scaffolding function, and may play a role in antiviral immunity. The associated SNP is predicted to be damaging to protein function using SIFT (Sorting Intolerant From Tolerant, a program that predicts whether an amino acid substitution affects protein function), but benign by Polyphen. rs2304256 is in the protein FERM (4.1 protein, ezrin, radixin, moesin) domain, which mediates interaction with Janus kinase and microtubule interacting protein 1, suggesting a possible increased interaction in DM cases. *TYK2* has been associated with rheumatoid arthritis,⁶ juvenile idiopathic arthritis,⁷ systemic lupus erythematosus,⁸ type 1 diabetes⁹ and multiple sclerosis.¹⁰ Two SNPs 2335 bp 5' of *BLK* (rs13277113) and within *BLK* (rs2618476) were associated with DM but not PM (table 2). These SNPs are highly correlated ($r^2 > 0.95$, HapMap CEU) with rs2736340, 7.5 kb 5' of *BLK*, supporting the previous DM GWAS results,¹ and suggesting a role of B cells in development of DM. No non-MHC SNPs were associated with PM at a Bonferroni corrected significance level.

We have identified *TYK2* as a novel associated locus for DM. This study confirms *TYK2* as an autoimmune gene and suggests a genetic overlap of DM with other autoimmune disorders, indicating genetic heterogeneity between PM and DM, although the difference may be due to the smaller PM sample size. These results require replication in an independent cohort and functional studies of the pathogenic role of *TYK2* in myositis. Our

analysis supports the study of additional immune-related loci in larger cohorts, using SNP arrays such as the ImmunoChip.

Acknowledgements

The authors thank the other members of the EUMYONET and Myositis Genetics Consortium (MYOGEN) for their scientific and logistic support. A full list of investigators who contributed to the UK MYONET (Adult Onset Myositis Immunogenetic Collaboration) and JDM study can be found in Miller FW *et al.*¹ This study also makes use of data generated by the Wellcome Trust Case-Control Consortium. A full list of the investigators who contributed to the generation of the data is available from <http://www.wtccc.org.uk>. Funding for the project was provided by the Wellcome Trust under award 076113.

Funding This work was funded by Arthritis Research UK grant 18474, in part by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences, the European Science Foundation for EuMyoNet support and Association Francaise Contre Les Myopathies (AFM). The UK JDM Cohort and Biomarker study was supported by grants from the Wellcome Trust UK (085860), Action Medical Research UK, (SP4252), The Myositis Support Group UK, Arthritis Research UK (14518) and The Henry Smith Charity. The JDM Cohort study is adopted onto the Comprehensive Research Network through the Medicines for Children Research Network (<http://www.mcrn.org.uk>). The Czech cohort was supported by MH CZ-DRO Institute of Rheumatology 00023728.

REFERENCES

1. Miller FW, Cooper RG, Vencovsky J, et al. Genome-wide association study of dermatomyositis reveals genetic overlap with other autoimmune disorders. *Arthritis Rheum.* 2013; 65:3239–47. [PubMed: 23983088]
2. Hindorff, LA.; MacArthur, J.; Morales, J.; European Bioinformatics Institute. et al. [accessed 21 Oct 2010] A Catalog of Published Genome-Wide Association Studies. <http://www.genome.gov/gwastudies>
3. Bohan A, Peter JB. Polymyositis and dermatomyositis (first of two parts). *N Engl J Med.* 1975; 292:344–7. [PubMed: 1090839]
4. Bohan A, Peter JB. Polymyositis and dermatomyositis (second of two parts). *N Engl J Med.* 1975; 292:403–7. [PubMed: 1089199]
5. Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet.* 2009; 5:e1000529. [PubMed: 19543373]
6. Eyre S, Bowes J, Diogo D, et al. High-density genetic mapping identifies new susceptibility loci for rheumatoid arthritis. *Nat Genet.* 2012; 44:1336–40. [PubMed: 23143596]
7. Hinks A, Cobb J, Marion MC, et al. Dense genotyping of immune-related disease regions identifies 14 new susceptibility loci for juvenile idiopathic arthritis. *Nat Genet.* 2013; 45:664–9. [PubMed: 23603761]
8. Suarez-Gestal M, Calaza M, Endreffy E, et al. Replication of recently identified systemic lupus erythematosus genetic associations: a case-control study. *Arthritis Res Ther.* 2009; 11:R69. [PubMed: 19442287]
9. Wallace C, Smyth DJ, Maisuria-Armer M, et al. The imprinted DLK1-MEG3 gene region on chromosome 14q32.2 alters susceptibility to type 1 diabetes. *Nat Genet.* 2010; 42:68–71. [PubMed: 19966805]
10. Mero IL, Lorentzen AR, Ban M, et al. A rare variant of the TYK2 gene is confirmed to be associated with multiple sclerosis. *Eur J Hum Genet.* 2010; 18:502–4. [PubMed: 19888296]

Table 1

Polymyositis, dermatomyositis and matched control sample numbers included in the final analysis

Country	DM cases			PM cases			Cases total	Controls total
	JDM	DM	DM total	JPM	PM	PM total		
1. Czech Republic	11	134	145	–	61	61	206	166
2. Hungary	14	71	85	2	79	81	166	90
3. Netherlands/Sweden	4	54	58	–	67	67	125	642
4. Spain	4	61	65	–	–	–	65	259
5. UK	183	203	386	6	195	201	587	2415
TOTAL	216	523	739	8	402	410	1149	3572

DM, dermatomyositis; JDM, juvenile dermatomyositis; JPM, juvenile polymyositis; PM, polymyositis.

Table 2

SNPs associated with dermatomyositis, polymyositis or IIM

Phenotype	Chr	SNP	Position (hg19)	Locus*	p Value	Risk/non-risk allele	RAF cases	RAF controls	β	p Heterogeneity	Disease association for non-HLA SNPs ³
DM	6	rs116152465	32603007	5' HLA-DQA1	1.48542E-05	G/A	0.25667	0.20135	0.40	0.31	
DM	19	rs2304256	10475652	TYK2	0.00020	C/A	0.75478	0.7067	-0.25	0.59	T1DM
DM	6	rs114424451	32413051	3' HLA-DRA1	0.00021	G/A	0.91272	0.86887	0.34	0.99	
DM	8	rs13277113	11349186	5' BLK	0.00045	A/G	0.28098	0.24403	0.25	0.54	SLE
DM	8	rs2618476	11352541	BLK	0.00062	C/T	0.28903	0.25312	0.24	0.56	SLE
DM	9	rs7020673	4291747	GLIS3	0.0023	C/G	0.53583	0.49292	-0.18	0.64	T1DM
DM	5	rs6859219	55438580	ANKRD55	0.0023	C/A	0.82304	0.79178	-0.24	0.80	RA
DM	2	rs13385731	33701890	RASGRP3	0.0049	T/C	0.947972	0.928019	-0.38	0.67	SLE
DM	12	rs11171739	56470625	5' ERBB3	0.0069	T/C	0.62421	0.57102	0.28	0.09	T1DM
DM	3	rs11712165	119118796	ARHGAP31	0.0071	G/T	0.40245	0.38066	0.16	0.85	Celiac
DM	2	rs10210302	234158839	5' ATG16L1	0.0094	C/T	0.51259	0.46989	-0.17	0.36	Crohn's
PM	6	rs116152465	32603007	5' HLA-DQA1	0.00044	G/A	0.33252	0.20135	0.76	0.01	
PM	6	rs114424451	32413051	3' HLA-DRA1	0.0033	G/A	0.910976	0.86887	0.35	0.53	
PM	6	rs615672	32574171	Intergenic (HLA-DRB1)	0.0035	C/G	0.51585	0.35732	0.52	0.01	
PM	14	rs4900384	98498951	Intergenic (c14orf64)	0.0072	A/G	0.67073	0.71415	0.23	0.74	T1DM
IIM	6	rs114424451	32413051	3' HLA-DRA1	9.09513E-06	G/A	0.912097	0.86887	0.35	0.88	
IIM	6	rs116152465	32603007	5' HLA-DQA1	1.04525E-05	G/A	0.28442	0.20135	0.51	0.06	
IIM	19	rs2304256	10475652	TYK2	0.00027	C/A	0.74672	0.7067	-0.21	0.51	T1DM
IIM	12	rs11171739	56470625	5' ERBB3	0.0054	T/C	0.62753	0.57102	0.30	0.02	T1DM
IIM	11	rs6590330	128311059	Intergenic (ETS1)	0.0064	A/G	0.12931	0.10279	0.22	0.65	SLE
IIM	17	rs7221109	38770286	Intergenic (SMARCE1)	0.0071	T/C	0.37882	0.34209	-0.15	0.44	T1DM
IIM	2	rs10210302	234158839	5' ATG16L1	0.0071	C/T	0.50503	0.46989	-0.14	0.45	Crohn's
IIM	8	rs13277113	11349186	5' BLK	0.0077	A/G	0.27101	0.24403	0.21	0.24	SLE
IIM	3	rs11712165	119118796	ARHGAP31	0.0080	G/T	0.39723	0.38066	0.14	0.80	Celiac
IIM	8	rs2618476	11352541	BLK	0.0082	C/T	0.279	0.25312	0.20	0.27	SLE

SNPs in bold reached Bonferroni significance.

* Gene is annotated where closest gene is <10 kb, otherwise annotated as intergenic with closest gene in parentheses. Chr, chromosome; DM, dermatomyositis; HLA, human leukocyte antigen; IIM, idiopathic inflammatory myopathy; PM, polymyositis; p value, p value from random-effects meta-analysis; p heterogeneity, p value of test for heterogeneity; β , effect size (META V.1.5); RA, rheumatoid arthritis; RAF, risk allele frequency; SLE, systemic lupus erythematosus; SNP, single nucleotide polymorphism; T1DM, type 1 diabetes mellitus.