

2018

# What's in a Name? Moving to Neuroscience-Based Nomenclature in Pediatric Psychopharmacology

R. S. Sultan

C. U. Correll

*Zucker School of Medicine at Hofstra/Northwell*

J. Zohar

G. Zalsman

J. Veenstra-VanderWeele

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/articles>



Part of the [Psychiatry Commons](#)

---

## Recommended Citation

Sultan RS, Correll CU, Zohar J, Zalsman G, Veenstra-VanderWeele J. What's in a Name? Moving to Neuroscience-Based Nomenclature in Pediatric Psychopharmacology. . 2018 Jan 01; 57(10):Article 4066 [ p.]. Available from: <https://academicworks.medicine.hofstra.edu/articles/4066>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact [academicworks@hofstra.edu](mailto:academicworks@hofstra.edu).



# HHS Public Access

Author manuscript

*J Am Acad Child Adolesc Psychiatry*. Author manuscript; available in PMC 2019 March 21.

Published in final edited form as:

*J Am Acad Child Adolesc Psychiatry*. 2018 October ; 57(10): 719–721. doi:10.1016/j.jaac.2018.05.024.

## What's in a Name? Moving to Neuroscience-based Nomenclature in Pediatric Psychopharmacology

**Ryan S Sultan, MD,**

Department of Psychiatry, Columbia University and the New York State Psychiatric Institute, New York, NY, USA.

**Christoph U Correll, MD,**

Department of Psychiatry, The Zucker Hillside Hospital, Northwell Health, Glen Oaks, NY, Hofstra Northwell School of Medicine, Hempstead NY and The Feinstein Institute for Medical Research, Manhasset, NY, USA and the Charité Universitätsmedizin, Department of Child and Adolescent Psychiatry, Berlin, Germany.

**Joseph Zohar, MD,**

Department of Psychiatry, Chaim Sheba Medical Center, Tel Hashomer, Israel.

**Gil Zalsman, MD,** and

Geha Mental Health Center and Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel.

Department of Psychiatry, Columbia University and the New York State Psychiatric Institute, New York, NY, USA.

**Jeremy Veenstra-VanderWeele, MD**

Department of Psychiatry, Columbia University and the New York State Psychiatric Institute, New York, NY, USA.

### Keywords

anxiety disorder autism; mood disorder; ADHD; psychotic disorder

---

Our traditional names for psychotropic medication classes lead to unnecessary confusion. As clinicians, we have grown comfortable with idiosyncratic names of psychotropic medications and forgotten how unclear and misleading they can be. For example, evidence shows that serotonin reuptake inhibitors help in pediatric anxiety disorders, but a parent with an anxious child might ask, “If you diagnosed my son with separation anxiety, why are you giving him an antidepressant?” Another parent might object to the use of a “stimulant” medication, “My daughter never slows down, the last thing she needs is a stimulant!” Similarly, an “antipsychotic” can be prescribed on-label to youth with mania, bipolar depression, tics, or irritability in autism but families and patients might be confused by or object to the implied label of being “psychotic”. Further, patients or family members may

---

All correspondence should be directed to Ryan Sultan, MD at the New York State Psychiatric Institute/Department of Psychiatry, College of Physicians and Surgeons of Columbia University, 1051 Riverside Drive Box 78, New York, NY (telephone: 646-774-5888, sultanr@nypsi.columbia.edu).

**Potential Conflicts of Interests and Financial Disclosures:** Dr. Sultan reports no biomedical conflicts of interest or financial disclosures.

not feel comfortable asking clarifying questions and simply do not return for follow up--- concluding that the provider does not understand their child.

The current psychotropic nomenclature is largely based on concepts and knowledge from the 1960s and 70s. In many cases, the first identified psychotropic effect became the preferred term. “Stimulants” promote wakefulness. “Antidepressants” improved mood. In other cases, names evolved until sticking on the most unique effect. “Major tranquilizers” became “antipsychotics”. Other class names, such as benzodiazepines and tricyclic “antidepressants”, were named based on their chemical structure. None of these terms reflect contemporary scientific knowledge of how these medicines act in the central nervous system. For example, “antidepressants” do not bind to ‘depression receptors’ to reverse them. Instead, they block monoamine transporters, with downstream effects. Further, noradrenergic ‘antidepressants’ are only effective in treating depression, while serotonergic reuptake inhibitor ‘antidepressants’ are effective for depression and OCD (Zohar et al., 2015, Locher et al., 2017). In fact, despite being labeled antidepressants, SSRIs and SNRIs, have a more favorable number needed to treat for disorders than depression (Jane Garland et al. 2016).

Beyond being out-of-date and inaccurate, the current psychotropic medication names adversely affect patient care. These names conflate diagnoses and targets of treatment. They contribute to stigma by aligning the medications (and patients) with specific disorders. “Antipsychotics” are a clear example. This class of medication has indications for depression, mania, aggression, irritability, tics, as well as psychosis (Pillay et al., 2017, Correll et al., 2011).

This issue is present beyond psychiatry. General medicine struggles with similar confusion. For example, beta-blockers are often considered anti-hypertensive medications despite having a litany of other indications such as heart failure, essential tremor, migraine prophylaxis, arrhythmias, and even anxiety. To address this, an internist treating heart failure would avoid describing a beta-blocker as an anti-hypertensive to prevent confusion that the medication is for blood pressure control. In psychiatry, the consequences of medication class misnomers are more problematic, particularly for children. Young children can be concrete in their understanding of medicines and may not be able to separate the name of the treatment from their diagnosis, believing they must be “depressed” if they are taking an “antidepressant”. Further, well-meaning parents may attempt to explain a medicine after the visit is complete, raising the risk that the name of the medicine’s class becomes a substitute for an appreciation of the target symptoms.

Our legacy naming is problematic from a clinician’s point of view as well. Generalized naming overshadows the distinctions between mechanisms of action. Under the current nomenclature, bupropion, mirtazapine, duloxetine, and fluoxetine are all “antidepressants”; however, they vary wildly in their mechanisms of actions, side effects, and indications. In particular, some “antidepressants,” such as serotonin reuptake inhibitors, are commonly used for anxiety; whereas others, such as bupropion, have no apparent benefit for anxiety. Further, when our naming fails to distinguish medicines, it is hard for patients to understand why one “antidepressant” should work after another has failed. Promoting a more nuanced

understanding for psychotropic medications provides better differentiation for both clinicians and patients when considering, initiating, and continuing treatments.

Concerned by the limitations of our current nomenclature in the context of our growing neuroscience knowledge, a group of international organizations created a Nomenclature Taskforce to develop an approach that is rooted in scientifically-grounded neuroscience. The taskforce designed a system using current scientific data to classify psychiatric medications by pharmacology-driven nomenclature, rather than by chemical structure or disease. Instead of grouping loosely related types of medicines under the non-specific title of “antidepressants”, they are named based on their pharmacology and mode of action. This naming provides useful information for the clinician and paves the way to improved understanding for patients. It also helps to reduce stigma and misunderstanding. Instead of “antidepressants” for enuresis, clinicians can talk about “recruiting brain systems to change the depth of your sleep”. Instead of giving “antipsychotics”, clinicians can discuss “targeting the dopamine system to help you with your tics”.

The resulting system, Neuroscience-based Nomenclature (NbN), includes more than 130 psychotropic medications and also includes several layers of information beyond basic pharmacology, such as clinical, regulatory, and neurobiological data designed to help providers make informed prescribing decisions (Table 1) (Zohar et al. 2015).

As part of the classification system, the NbN expands and replaces our terminology into eleven pharmacological domains, such as Norepinephrine, Dopamine, Glutamate, and Serotonin. Ten modes of action are also identified, ranging from effects at receptors and transporters to impacts on ion channels and enzymes. These pharmacological domains and modes of action are cornerstones for a nuanced description of psychotropic medications.

Despite the many advantages to the NbN terminology, challenges exist to adopting this classification system. A primary challenge is the inertia we have as clinicians who have used the legacy nomenclature for our whole careers. However, pushing past this inertia has significant benefits. For example, when providing psychoeducation about ADHD, explaining the etiology of ADHD in the context of the dopamine and norepinephrine systems logically flows into a discussion of medication treatment---methylphenidate and amphetamine salts as dopamine and norepinephrine reuptake inhibitors and releasers. Dropping the term “stimulant” and talking about the mechanism of action creates a more insightful, coherent and less misleading explanation.

Beyond the inherent barrier of changing how we talk about medicines, another challenge is that our understanding of medication mechanisms and molecular pharmacology remains incomplete. This issue is most clearly evidenced by lithium, which currently sits in a category of its own because its exact mechanism remains uncertain despite established actions on various enzymatic targets (Mota de Freitas et al., 2016). However, despite this challenge, NbN is an improvement over lumping lithium with valproic acid and lamotrigine as “mood stabilizers”, as if their pharmacology had anything in common.

As part of NbN’s goal to be current and accessible, a free app (NbN) is available for common mobile device platforms--- including full descriptions of psychotropic medications,

their former terminology, pharmacology targets, modes of action, approved indications, efficacy (off-label use), side effects, practical notes and neurobiology. A separate child psychiatry specific app (NbN-ca) has also been developed. Different from the adult NbN app, it includes specific child-based dosing information as part of the practical notes. It also includes data on age of approval by major regulatory agencies. Through these apps, the NbN is a living document that can easily be updated and improved as new information becomes available.

The adoption of neuroscience-based nomenclature represents an opportunity for growth and improvement in our field's descriptions of the primary biological treatment when using psychopharmacology. This system offers a less stigmatizing, more scientifically grounded, easily updatable, and nuanced understanding of our treatments. Concurrently, it exposes the complexity and limitations of the field's knowledge of psychopharmacology. However, this problem is also an opportunity for improved education of our patients and ourselves. We can use these new terms to empower youth and families to better understand what their treatments do, without a distracting legacy of misnomers.

## Acknowledgments

Dr. Correll has received grant or research support from the National Institute of Mental Health, the Patient Centered Outcomes Research Institute, The Bendheim Foundation, Janssen and Takeda. He has served as a member of advisory boards/the Data Safety Monitoring Boards for Alkermes, IntraCellular Therapies, Lundbeck, Neurocrine, Otsuka, Pfizer, and Sunovion. He has served as a consultant to Alkermes, Allergan, the Gerson Lehrman Group, IntraCellular Therapies, Janssen/Johnson & Johnson, LB Pharma, Lundbeck, Medscape, Otsuka, Pfizer, ProPhase, Sunovion, Supernus, and Takeda. He has presented expert testimony for Bristol-Myers Squibb, Janssen, and Otsuka. He has received honoraria from Medscape. He has received travel expenses from Janssen/Johnson and Johnson, Lundbeck, Otsuka, Pfizer, Sunovion, and Takeda.

Dr. Veenstra-VanderWeele has received grant or research support from the National Institute of Mental Health, the National Center for Child Health and Human Development, the National Institute for Neurological Disease and Stroke, the Agency for Healthcare Research and Quality, the Health Resources and Services Agency, the Simons Foundation Autism Research Institute, Autism Speaks, Roche, Novartis, Forest, Seaside Therapeutics, and SynapDx. He has served on advisory boards for Autism Speaks, the Brain and Behavior Research Foundation, Roche, Novartis, and SynapDx. He has received editorial stipends from Springer and Wiley. He has received travel expenses from Roche, Novartis, and SynapDx.

Dr. Zohar has received grant/research support from Lundbeck, Servier, Brainsway and Pfizer, has served as a consultant or on advisory boards for Servier, Pfizer, Abbott, Lilly, Actelion, AstraZeneca and Roche, and has served on speakers' bureaus for Lundbeck, Roch, Lilly, Servier, Pfizer and Abbott.

Dr. Zalsman has served as a consultant for Janssen and Elminda.

## References:

- Mota de Freitas D, Levenson BD, Goossens JL (2016) Lithium in Medicine: Mechanisms of Action In: Sigel A, Sigel H, Sigel R (eds) *The Alkali Metal Ions: Their Role for Life. Metal Ions in Life Sciences*, vol 16 Springer, Cham
- Zohar J, Stahl S, Moller HJ, et al. (2015). A review of the current nomenclature for psychotropic agents and an introduction to the neuroscience-based nomenclature. *Euro Neuropsychopharmacol.* 25:2318–2325.
- Locher C, Koechlin H, Zion SR, et al. Efficacy and Safety of Selective Serotonin Reuptake Inhibitors, Serotonin-Norepinephrine Reuptake Inhibitors, and Placebo for Common Psychiatric Disorders Among Children and Adolescents: A Systematic Review and Meta-analysis. *JAMA Psychiatry.* 2017 10 1;74(10):1011–1020. [PubMed: 28854296]

- Pillay J, Boylan K, Carrey N, et al. First- and Second-Generation Antipsychotics in Children and Young Adults: Systematic Review Update [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2017 Mar. Available from <https://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0096139/>. Accessed April 1, 2018
- Correll CU, Kratochvil CJ, March JS. Developments in pediatric psychopharmacology: focus on stimulants, antidepressants, and antipsychotics. *J Clin Psychiatry*. 2011 5;72(5):655–70. [PubMed: 21658348]
- Jane Garland E, Kutcher S, Virani A. et al. Update on the Use of SSRIs and SNRIs with Children and Adolescents in Clinical Practice. *J Can Acad Child Adolesc Psychiatry*. 2016 Winter; 25(1): 4–10. Published online 2016 Feb 1. PMID: PMC4791100 PMID: 27047551 [PubMed: 27047551]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 1:**

Layers of information in NbN system

Characteristic		Example: Methylphenidate
1	Mode of Action	Reuptake inhibitor (DAT, NET), releaser (DA, NE) New Terminology
2	Pharmacology Domain	Dopamine, Norepinephrine
3	Approved indications	Attention deficit hyperactivity disorder in children >6 years old and adults (FDA, EMA)
4	Efficacy	Reduces signs and symptoms of attention deficit hyperactivity disorder in adults and children; Used to treat narcolepsy
5	Side Effects	Headache, insomnia, nervousness, decreased appetite.
6	Practical notes	(d) enantiomer used less often than the racemic mixture. A number of slow release formulations are available with longer durations of action (usually once daily) and lower abuse liability
7	Former Terminology	Stimulant
8	Neurobiology	<p><b>Neurotransmitter effects preclinical</b> Blocks DA transporter and to a lesser extent NE transporter</p> <p><b>Physiological human</b> Promotes wakefulness, increased blood pressure and heart rate, insomnia. Increased ratings of "active/alert/energetic", "stimulated," "shaky," and "jittery"</p> <p><b>Brain circuits human</b> Normalizes abnormal cognitive function and associated fMRI signals in patients with attention deficit hyperactivity disorder</p>

Source: NBN Website (<https://www.nbn2.com/search#drug/70/rid/70>)