

2018

GRK5 is associated with the regulation of methadone dosage in heroin dependence

S. C. Wang

R. H. Chung

H. W. Kuo

T. H. Liu

C. P. Fang

*See next page for additional authors*Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/articles>Part of the [Psychiatry Commons](#)

Recommended Citation

Wang SC, Chung RH, Kuo HW, Liu TH, Fang CP, Liu SC, Liu CC, Tsou HH, Chen AC, Liu YL. GRK5 is associated with the regulation of methadone dosage in heroin dependence. . 2018 Jan 01; 21(10):Article 4072 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/articles/4072>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

Authors

S. C. Wang, R. H. Chung, H. W. Kuo, T. H. Liu, C. P. Fang, S. C. Liu, C. C. Liu, H. H. Tsou, A. C. Chen, and Y. L. Liu

REGULAR RESEARCH ARTICLE

GRK5 Is Associated with the Regulation of Methadone Dosage in Heroin Dependence

Sheng-Chang Wang, Ren-Hua Chung, Hsiang-Wei Kuo, Tung-Hsia Liu, Chiu-Ping Fang, Shu Chih Liu, Chia-Chen Liu, Hsiao-Hui Tsou, Andrew C. H. Chen, Yu-Li Liu

Center for Neuropsychiatric Research, National Health Research Institutes, Miaoli County, Taiwan (Dr Wang, Ms Kuo, Ms T.-H. Liu, Ms Fang, Ms S.C. Liu, Ms C.-C. Liu, and Dr Y.-L. Liu); Division of Biostatistics and Bioinformatics, Institute of Population Health Sciences, National Health Research Institutes, Miaoli County, Taiwan (Drs Chung and Tsou); Graduate Institute of Biostatistics, College of Public Health, China Medical University, Taichung, Taiwan (Dr Tsou); Department of Psychiatry, the Zucker Hillside Hospital, Northwell Health, Glen Oaks, New York (Dr Chen); The Feinstein Institute for Medical Research, Hofstra Northwell School of Medicine at Hofstra University, Manhasset, New York (Dr Chen); Graduate Institute of Clinical Medical Science, China Medical University, Taichung, Taiwan (Dr Y.-L. Liu).

Correspondence: Yu-Li Liu, PhD, Center for Neuropsychiatric Research, National Health Research Institutes, 35 Keyan Road, Zhunan, Miaoli County 350, Taiwan (yliou@nhri.org.tw).

Abstract

Background: There is no countable biomarker for opioid dependence treatment responses thus far. In this study, we recruited Taiwanese methadone maintenance treatment patients to search for genes involving the regulatory mechanisms of methadone dose by genome-wide association analyses.

Methods: A total of 344 Taiwanese methadone maintenance treatment patients were included in a genome-wide association study. The involvement of GRK5 in opioid dependence was then further confirmed by gene expression study on lymphoblastoid cell lines derived from 3 independent age- and gender-matched groups: methadone maintenance treatment patients, medication-free former heroin abusers, and normal controls.

Results: The results indicated that GRK5, the gene encoding an enzyme related to μ -opioid receptor desensitization, is associated with methadone dose by additive model of gene-based association analysis ($P = 6.76 \times 10^{-5}$). We found that 6 of the 55 single nucleotide polymorphisms from the genome-wide genotype platform and 2 single nucleotide polymorphisms from the 29 additionally selected single nucleotide polymorphisms were significantly associated with methadone maintenance dose in both genotype and allele type ($P \leq .006$), especially in patients who tested negative in the urine morphine test. The levels of GRK5 gene expression were similar between methadone maintenance treatment patients and medication-free former heroin abusers. However, the normal controls showed a significantly lower level of GRK5 gene expression than the other groups ($P = .019$).

Conclusions: The results suggested an important role for GRK5 in the regulatory mechanisms of methadone dose and course of heroin dependence.

Keywords: GRK5, haplotype, methadone maintenance dose, MMT, SNP

Received: March 2, 2018; Revised: June 24, 2018; Accepted: July 25, 2018

© The Author(s) 2018. Published by Oxford University Press on behalf of CINP.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Significance Statement

There is no countable biomarker for opioid dependence and its treatment responses thus far. In this study, we first recruited 344 Taiwanese methadone maintenance treatment (MMT) patients to search for genes involved in the regulatory mechanisms of methadone dose by genome-wide association analyses. The results indicated that *GRK5*, the gene encoding an enzyme related to μ -opioid receptor desensitization, is associated with methadone dose. The involvement of *GRK5* in opioid dependence was then further confirmed by gene expression study on cell lines derived from 3 independent age- and gender-matched groups: MMT patients, medication-free former heroin abusers, and normal controls. Our findings provide novel evidence suggesting that *GRK5* could be a useful indicator for the severity of opioid dependence and a biomarker for methadone dosage. This study is the first clinical research supporting *GRK5* as a candidate gene involving the regulation of methadone dosage in opioid dependence.

Introduction

Methadone is a well-established medication for treatment of heroin dependence (Mattick et al., 2009). Methadone maintenance therapy (MMT) has shown superior treatment effectiveness particularly in terms of patient retention rate and reduction in heroin use compared with no treatment (Mattick et al., 2002, 2003, 2009). However, determination of an optimal methadone dosage remains a challenge for clinicians because of several serious potential adverse effects. The optimal methadone dosage is affected by various physiological and psychological statuses of patients (Hiltunen and Eklund, 2002; Murray et al., 2008; Sullivan et al., 2014; Mouly et al., 2015). A steady-state methadone concentration in the blood is usually achieved when patients receive daily treatment of methadone for longer than 5 times the half-life (Chan and Sim, 2015). A steady-state dosage is influenced by both individual metabolic activity and receptor-coupled mechanisms of actions toward the treatment responses. Thus, it would be a better sample to use MMT patients who had reached a steady dosage of treatment in search of genes related to dosage regulation.

A few genes encoding drug transporters, P-glycoprotein transporters, have been reported to be associated with methadone dose, for example, *ABCB1* (ATP binding cassette subfamily B member 1) gene in a Jewish (Levrant et al., 2008) and an Australian cohort (Coller et al., 2006) respectively. The gene encoding liver metabolic enzyme cytochrome P-450 (CYP) 2B6 has also been reported to be associated with methadone dose in an Israeli study (Levrant et al., 2013a). Most studies reported a set of genes that are involved with both metabolic and receptor coupled pathways. For example, *ABCB1* and μ -opioid receptor *OPRM1* genes in Australian methadone MMT patients (Barratt et al., 2012); CYPs and *ABCB1* genes in Caucasian patients (Fonseca et al., 2011); *BDNF*, *NTRK2*, *OPRM1*, *DRD2*, and *ANKK1* genes in Israeli patients (Levrant et al., 2013b); and *ABCB1*, *CYP2B6*, *OPRM1*, *ANKK1*, and *DRD2* genes in Taiwanese patients (Hung et al., 2011). These reports suggest that more than one single gene is involved with the biological mechanisms underlying the regulation of methadone dosage.

In our previous studies in a Taiwanese MMT patient cohort, we found that *CYP2C19* (Wang et al., 2013b), *2B6*, *2C19*, *3A4* (Tsai et al., 2014), and *OPRM1* genes (Wang et al., 2012) had demonstrated marginally significant associations with methadone dose. In the present study, we performed a genome-wide association study with 344 MMT patients and analyzed genes associated with steady-state methadone dose. The gene encoding guanine nucleotide-binding protein (G protein)-coupled receptor kinase 5 (*GRK5*) was identified as a gene significantly associated with methadone dosage. Hence, more SNPs in the significant genetic regions in *GRK5* were further selected and genotyped. *GRK5* is an enzyme that belongs to the G protein-coupled receptor kinase

subfamily of the Ser/Thr protein kinase family (Pitcher et al., 1998). It has been reported that *GRK5* is responsible for desensitization of μ -opioid receptor (Mann et al., 2015). We speculate that this may be one of the mechanisms through which methadone dosage is regulated.

Methods

All methods in this study were performed in accordance with the guidelines and regulations of the institutional review committees. The study protocol was approved by the institutional review boards of the National Health Research Institutes (EC0970504, Zhunan, Taiwan) and the 7 participating hospitals: Tao-Yuan Mental Hospital, En-Chu-Kong Hospital, Far-Eastern Memorial Hospital, Taipei City Hospital Song-De and Yang-Ming Branches, China Medical University Hospital, and Wei-Gong Memorial Hospital (Wang et al., 2013a). Written informed consent was obtained from all participants. The project has also been registered with the National Institutes of Health (NIH) Clinical Trial (<https://clinicaltrials.gov/ct2/show/NCT01059747> for 344 MMT patients). The inclusion criteria were aged ≥ 18 years, receipt of MMT for at least 3 months with regular attendance for the past 7 days, and a methadone dosage adjustment of no more than 10 mg in the past 7 days. Exclusion criteria were co-morbidity with medical or other mental disorders requiring immediate treatment and pregnancy.

The genome-wide genotype data were obtained from a cross-sectional study recruited from year 2008 to 2009. A total of 344 MMT patients had passed the genome-wide association study (GWAS) quality controls with available detailed data regarding plasma methadone concentrations and urine morphine test results (Yang et al., 2016). To further verify the role of *GRK5* in heroin addiction and methadone treatment, additional age- and gender-matched normal controls ($n=23$) who did not have any psychiatric or substance abuse history, and former heroin abusers ($n=23$) who were medication-free and abstinent for average 2 years were included in the gene expression study. The gene expression study protocol was approved by the institutional review board of the National Health Research Institutes (EC0980209-R5, Zhunan, Taiwan). Written informed consents were also obtained from all participants. The former heroin abusers were recruited mainly from nongovernment organization-affiliated, medication-free therapeutic communities. Subjects in the medication-free former-heroin abuser group met the following criteria: (1) lifetime DSM-IV diagnosis of opioid dependence; and (2) no illicit drug use other than opioid or alcohol intoxication in the last 4 weeks. The normal control subjects were recruited from referrals, advertisements, and posted notice, and they met the following criteria: (1) no lifetime

DSM-IV diagnosis of opioid abuse or dependence; and (2) no any illicit drug use nor alcohol intoxication in the past 28 days (<https://clinicaltrials.gov/ct2/show/results/NCT01668706> for former heroin abusers and normal controls of NIH registration). All these subjects were included for comparison in the gene expression study. Urine specimens were collected for screening of illicit drug use. Venous blood was taken for routine biochemical examinations and viral infectious disease profile.

Clinical Assessments

Demographics, clinical characteristics, and methadone treatment courses, including the methadone dose, treatment duration, and treatment adherence in the previous week, were obtained from medical records. Several interviewer-administered assessments including the Treatment Outcomes Profile (Marsden et al., 2008), which measures the amount and frequency of alcohol and other illicit substance use in the past 28 days, and the Clinical Opioid Withdrawal Scale (COWS), which measures the severity of 11 opioid withdrawal symptoms (Wesson and Ling, 2003), were conducted before methadone was administered on the test day. A COWS score >5 is considered as significant withdrawal symptoms (Tompkins et al., 2009). Anti-hepatitis C virus (HCV) antibody was measured by electrochemiluminescence immunoassay on Cobas e601 (Roche Diagnostics, Basel, Switzerland), and anti-HIV antibody was measured using Abbott chemiluminescence immunoassay (Abbott Diagnostics, Chicago, IL). The levels of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were measured with Kinetic UV method. The level of gamma-glutamyl transpeptidase (γ -GT) was measured with enzymatic colorimetric assay. These assays were performed using the Integra 800 device (Roche Diagnostics).

Urine Morphine Test

Urine specimens were collected prior to administration of methadone on the study day. In our present analyses and previous reports, the urine morphine test was used as a surrogate measurement for the methadone treatment outcome.

Genome-Wide SNP Genotyping

Genomic DNA was extracted from the buffy coat of 6 mL whole blood. Genotyping was performed with the Axiom Genome-Wide CHB 1 Array, which was population-optimized to have a better genomic coverage of common alleles (minor allele frequency >5%) of the Han Chinese genome. All sample call rates were >98.8%, and the mean individual sample call rate was $99.7 \pm 0.18\%$. Details of quality control and raw data can be accessed in Gene Expression Omnibus (GEO accession number: GSE78098; see also supplementary Methods) (Yang et al., 2016).

GRK5 SNP Selection and Genotyping

Genotypes of the additionally selected 29 SNPs in GRK5 were identified by matrix-assisted laser desorption/ionization-time of flight mass spectrometry (Rodi et al., 2002). In brief, the primers and probes flanking the SNPs were designed using SpectroDESIGNER software (Sequenom, San Diego, CA). DNA fragments (100–300 bp) encompassing each SNP site were amplified by PCR (GeneAmp 9700 thermocycler, Applied Biosystems, Foster City, CA) in accordance with the manufacturer's instructions. After the unincorporated dNTPs were removed and the

PCR reaction was inactivated by shrimp alkaline phosphatase, primer extension was performed by adding appropriate probes to the PCR reaction. Various nucleotide extension products were differentiated by matrix-assisted laser desorption/ionization-time of flight analysis (Tost and Gut, 2005).

EBV-Transformed Lymphoblastoid Cell Culture

Lymphocytes from subjects were transfected with Epstein-Barr virus (EBV) then cultured as lymphoblastoid cell line for gene expression assay (see also supplementary Methods). Total RNA of EBV-transformed lymphoblastoid cells was extracted by TRIzol Reagent (Life Technologies, Inc., Gaithersburg, MD).

GRK5 Gene Expression

A total of 63 MMT patients selected from the 344 MMT patients in consideration of minor genotype carriers, 23 age- and gender-matched normal controls, and 23 medication-free abstinent former heroin abusers were included in the gene expression study (see also supplementary Methods). Gene expression was quantified relatively to TBP expression, the relative quantification method, using ABI StepOne Plus Software. The relative expression level of GRK5 compared with that of TBP was defined as $-\Delta\Delta CT = -[CT_{GRK5} - CT_{TBP}]$, where CT was the cycle threshold. The GRK5 mRNA/TBP mRNA ratio was calculated from $2^{-\Delta\Delta CT} \times K$, in which K was a constant.

Statistical Analyses

Summary statistics such as means and SDs were used to describe the samples. Means among different cohorts were compared using the Kruskal-Wallis test for 3-group comparison or Mann-Whitney U-test for 2-group. Categorical variables among cohorts were compared using a standard chi-square test. Spearman correlation coefficient was used to test the associations between the methadone dosage and total COWS score. The aforementioned analyses were conducted using SAS software, Version 9.4 (SAS Institute, Inc., Cary, NC).

Gene-based analyses were performed using the Knowledge-based mining system for Genome-wide Genetic studies (KGG) (Li et al., 2012). Single-SNP association P values were first calculated using PLINK (Purcell et al., 2007). In KGG, the SNPs were divided into linkage disequilibrium blocks, and a score based on the extended Sime's test (Li et al., 2011) was calculated for each block using the single-SNP P values. The scores for blocks within a gene were combined into a scaled chi-square statistic as a hybrid set-based test. KGG also produced Q-Q plots for the gene-based P values, single-SNP P values for SNPs in genes, and single-SNP P values for SNPs outside genes. We then calculated the genomic inflation factor, lambda value (Devlin and Roeder, 1999), for the gene-based P values.

To further evaluate the effects of SNPs in a candidate gene identified by KGG, single SNP association tests were performed for SNPs that were more densely genotyped in that gene. The genotypic or allelic models for testing associations of SNPs with the treatment outcomes were performed using the Correlation/Trend test provided in SNP & Variation Suite, Version 8.4.0 (Golden Helix, Inc., Bozeman, MT). We also performed haplotype association tests in these densely genotyped SNPs. Global P values of the haplotype association tests for methadone dose were calculated using the generalized estimating equation model. The haplotype analysis was also conducted using SAS software, Version 9.4 (SAS Institute, Inc). Mann-Whitney U tests were

performed to compare the difference in gene expression levels between groups. The 2D dot plot and the Manhattan plot, which presents the $-\log_{10}$ of genome-wide P values, were calculated and plotted by GraphPad Prism 5 (GraphPad Software, San Diego, CA). The regional visualization of association plots was plotted by LocusZoom (Pruim et al., 2010).

False discovery rate was employed to correct for multiple comparisons, that is, an adjusted P value is defined as the smallest significance level for which the given hypothesis would be rejected when the entire family of tests is considered (Storey, 2003). Hardy-Weinberg equilibrium tests were performed using HAPLOVIEW version 4.2 (Barrett et al., 2005). A statistically significant level was based on a 2-sided test and assigned for $P < .05$.

RESULTS

Clinical Characteristics and Methadone Treatment Outcomes

A total of 344 MMT patients passed the quality controls (Yang et al., 2016) with an average age around 38 years, and the percentage of males was 82 (Table 1). All patients had taken methadone daily in the previous 1 week. The average total withdrawal symptom scores rated by COWS >5 ($n=23$) were 6.6 ± 2.3 in the 344 MMT patients. The methadone dose did not show correlations with the total COWS score >5 (Spearman correlation coefficient, $n=23$, $r=0.34$, $P=.12$ in MMT patients). The clinical characteristics and single gene observations from this cohort were well characterized for the recruitment and treatment responses in our previous studies (Wang et al., 2013a).

Identification of Genes Associated with the Methadone Dose Using Gene-Based Models in the GWAS

PLINK linear regression was employed for the 344 subjects and 615216 SNPs. KGG was employed for the gene-based model analyses. Total of 15553 genes were tested for associations with methadone dose in the gene-based tests using the additive model. The methadone dose was most statistically significant associated with the GRK5 gene in the gene-based test using the additive test (supplementary Figure 1A; supplementary Table 1). The genomic inflation factor, lambda value, was 0.892,

Table 1. Demography of the MMT Patients

Variable	n	Mean \pm SD
Age	344	38.16 \pm 7.69
Gender [male (%)]	281	(81.69)
BMI (kg/m ²)	341	23.64 \pm 3.52
Methadone dosage (mg/d)	344	55.22 \pm 28.47
Sum of COWS	344	1.51 \pm 1.87
Urine morphine (+) (%)	173	(50.58)
HCV (+) (%)	313	(94.56)
HIV (+) (%)	76	(22.55)
Liver function test		
AST (U/L)	326	52.42 \pm 57.06
ALT (U/L)	332	61.13 \pm 75.30
γ -GT (U/L)	310	63.38 \pm 104.68

Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase; COWS, Clinical Opioid Withdrawal Scale; MMT, methadone maintenance treatment; γ -GT, gamma-glutamyl transpeptidase.

suggesting that there were no population stratification and systematic genotyping errors in our sample. The Q-Q plots in supplementary Figure 1B showed that the P values for gene-based tests and for SNPs within and outside genes mostly fell within the 95% CIs of the expected values, again suggesting that there was no inflation of type I error rates in our tests.

SNPs in GRK5 Were Associated with the Methadone Maintenance Dose

GRK5 gene is located at the 10q26.11 chromosome region, spans approximately 248 kb pairs, and contains 16 exons. A total of 55 SNPs within the GRK5 genetic region were placed in the genome-wide genotyping platform in which 6 (rs4752269, rs2061078, rs12415832, rs4752300, rs10787966, rs1889432) of the 55 SNPs (supplementary Table 2) showed significant associations with methadone dose. To confirm the genetic association results with methadone dose, 29 additional SNPs were selected according to the minor allele frequency >0.1 on the HapMap of Chinese ethnic group (<http://hapmap.ncbi.nlm.nih.gov>), the functions predicted by the bioinformatics FastSNP (Yuan et al., 2006), and reports in the literature (Jakobsdottir et al., 2005; Arawaka et al., 2006; Tarantino et al., 2011). Six (rs871196, rs11819686, rs12780837, rs10886472, rs11198907, and rs1537576) of the 29 SNPs showed significant associations with methadone dose. All SNPs were in Hardy-Weinberg equilibrium. Eight SNPs (rs4752269, rs2061078, rs12415832, rs10886472, rs11198907, rs4752300, rs10787966, and rs1889432) were significantly associated with methadone dose in MMT patients (Figure 1). The minor allele type carriers of these 8 SNPs had a higher methadone dose than the major allele type carriers in all MMT patients. SNP rs10886472 showed the highest level of significance in association with the methadone dose in all MMT patients (supplementary Table 3). The significant associations were mainly contributed from the urine morphine test-negative MMT patients (supplementary Table 4). In addition, it remains statistically significant after adjusting for HCV infection rate. The allele frequencies of these SNPs are similar to those in the Han Chinese in Beijing, China population in the National Center for Biotechnology Information (supplementary Table 5).

GRK5 Haplotypes Were Associated with Methadone Dose

Two haplotype blocks were created by Haploview from the 8 SNPs in GRK5 (supplementary Figure 2). They are block 1 located at the intron 3–4 consisting of 3 SNPs (rs10886472-rs11198907-rs4752300), and block 2 located at the intron 4–6 consisting of 2 SNPs (rs10787966-rs1889432) (supplementary Table 6A). The major haplotypes (CGG haplotype frequency 76% in block 1, and GA haplotype frequency 77% in block 2) within each block were significantly associated with methadone dose. Carriers with the major haplotype had a lower methadone dose than those with the minor haplotype. The haplotype blocks that showed significant associations with methadone dose were mainly observed in the urine morphine test-negative patients (supplementary Table 6B).

Functional SNP Estimation in GRK5

rs11198907, rs4752300, and rs10787966 in the intron 4, among the 8 SNPs, showed significant associations with the level of GRK5 gene expression in the genotype recessive model (supplementary Table 7). The significant level of associations was higher




Figure 1. Correlation/trend association of the 8 single nucleotide polymorphisms (SNPs) in guanine nucleotide-binding protein (G protein)-coupled receptor kinase 5 (GRK5) gene-differentiated treatment response according to methadone dosage in 344 methadone maintenance treatment (MMT) patients using the regional visualization of association plots in LocusZoom.

in the MMT patients who tested negative in the urine morphine test (Mann-Whitney U test, $P = .048$, $.031$, and $.041$, respectively, in all patients vs $.021$, $.021$, and $.035$, respectively, in the MMT patients with a negative urine morphine test) (supplementary Figure 3).

MMT Patients and Medication-Free Former Heroin Abusers Showed Higher Level of GRK5 Gene Expression Than Normal Controls

To further verify the role of GRK5 gene in opioid dependence, we recruited additional former heroin abusers ($n = 23$) who had been medication-free and abstinent from opioid use for at least 2 years and normal controls ($n = 23$) who had never used any illicit substance. They were compared with 63 age- and gender-matched subjects from the 344 MMT patients (Table 2). MMT patients showed a similar level of GRK5 gene expression compared with medication-free former heroin users. However, MMT patients showed a significantly higher level of GRK5 gene expression in culture lymphoblastoids than normal controls (Mann-Whitney U test, $P = .019$) (Figure 2).

Discussion

The present study was designed to identify genes associated with methadone treatment dose for opioid dependence by GWAS. GRK5 gene expression assays were further tested in medication-free abstinent former heroin abusers, normal controls, and patients under MMT to confirm the involvement of GRK5 in the course of opioid addiction and pharmacodynamics of methadone.

The activity of mu-opioid receptor in processing consequent signal transduction is regulated by several G protein-coupled receptor kinases (Mann et al., 2015). Phosphorylation of the mu-opioid receptor by GRK5 could initiate further hierarchical phosphorylation cascade. The results in this study support the role of GRK5 in the regulatory mechanism of methadone dosage. We found in the current GWAS study that a genetic polymorphism, rs10886472 (intron3), within the 8 SNPs located from introns 1 to

6 of GRK5 gene showed the highest significant level in association with methadone maintenance dose. The significant associations were mainly contributed from the urine morphine test-negative MMT patients, which represent a subgroup of patients who had a better methadone treatment outcome. On the contrary, association between GRK5 and methadone dose was not found in those MMT patients with a positive urine morphine test. The haplotype blocks constructed by the 8 SNPs in GRK5 also demonstrated strong significant associations with methadone dose. In the GRK5 gene expression tendency analyses of rs11198907, rs4752300, and rs10787966 located within intron 4 (supplementary Figure 3), major genotype carriers showed a higher level of GRK5 expression and a lower methadone dose than the minor genotype carriers. In an animal study, the *Grk5* knockout mice lost the manifestation of morphine reward and dependence (Glück et al., 2014). These observations all suggest crucial roles of GRK5 in the regulation of methadone dosage and course of heroin addiction. The possible effects of GRK5 gene expression on the decreased reward sensation in the heroin-addictive patients under MMT warrant further investigation.

A recent genome-wide association study where the methadone doses were obtained by self-report from subjects identified a new significant locus located at the upstream of OPRM1 gene (Smith et al., 2017). However, the most significant SNP, rs73568641, in the report showed a very low minor allele frequency (0.04) in Han Chinese in Beijing, China and JPT populations (https://www.ncbi.nlm.nih.gov/projects/SNP/snp_ref.cgi?rs=73568641). We therefore examined a few other SNPs located nearby rs73568641 and found that OPRM1 genetic loci showed only marginal significant associations with methadone dose in our subjects where the most significant P value was $.004$ in single SNP association analysis (supplementary Figure 4).

Elevated level of GRK5 gene expression in the lymphoblastoids of MMT patients and medication-free former heroin abusers may be related to desensitization of mu-opioid receptor, which may indicate a more severe status of chronic opioid dependence. In this study, GRK5 gene expression among the 3 age- and gender-matched cohorts showed a rank order of expression level as follows: The levels of gene expression are

Table 2. General Demography of the Subjects Recruited for the Verification Study: Patients under Methadone Treatment (MMT), Medication-Free Former Heroin Abusers, and Healthy Normal Controls

Variable	MMT patients (n=63)		Former heroin abusers (n=23)		Normal Controls (n=23)		P value
	n	Mean ± SD	n	Mean ± SD	n	Mean ± SD	
Age	63	38.14 ± 6.13	23	38.04 ± 6.73	23	37.39 ± 6.65	.876 ^a
Gender							.763 ^b
Male	50	(79.37%)	19	(82.61%)	17	(73.91%)	
Female	13	(20.63%)	4	(17.39%)	6	(26.09%)	
BMI	63	23.91 ± 3.36	22	25.98 ± 3.25	23	24.17 ± 4.05	.012 ^a
Cotinine (ng/mL)	63	350.72 ± 169.32	23	13.32 ± 48.35	23	39.69 ± 81.03	<.001 ^a
Nonsmoking	2	(3.17%)	19	(82.61%)	16	(69.57%)	<.001 ^b
Smoking	61	(96.83%)	4	(17.39%)	7	(30.43%)	
HIV							.021 ^b
Negative	47	(77.05%)	20	(90.91%)	23	(100.00%)	
Positive	14	(22.95%)	2	(9.09%)	0	(0.00%)	
HCV							<.001 ^b
Negative	5	(8.62%)	8	(36.36%)	23	(100.00%)	
Positive	53	(91.38%)	14	(63.64%)	0	(0.00%)	
Liver function test							
AST (U/L)	60	45.05 ± 34.13	22	48.91 ± 38.94	23	25.48 ± 10.24	.001 ^a
ALT (U/L)	61	54.87 ± 48.55	22	77.55 ± 76.34	23	30.78 ± 20.44	.012 ^a
γ-GT (U/L)	59	51.88 ± 95.77	22	40 ± 25.13	23	55.13 ± 63.62	.825 ^a

Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase; γ-GT, gamma-glutamyl transpeptidase; HCV, Hepatitis C virus antibody; HIV, Human Immunodeficiency Virus.

^a Kruskal-Wallis test.

^b Chi-square test. P values < .05 were shown in bold font.




Figure 2. Levels of guanine nucleotide-binding protein (G protein)-coupled receptor kinase 5 (GRK5) gene expression among methadone maintenance treatment (MMT) patients, medication-free abstinent former heroin abusers (abstinent), and normal controls (normal). Data were presented as means ± SD. N: number of subject.

similar between MMT patients and medication-free abstinent former heroin abusers, but lower in the normal controls group. To our knowledge, our result is the first human study suggesting that MMT patients may potentially show a higher level of desensitization of mu-opioid receptor, as shown in the level of GRK5 gene expression. As mentioned above, phosphorylation of receptor is one of the mechanisms through which GRK5 influences desensitization of mu-opioid receptor (Mann et al., 2015).

The results also support the hypothesis that GRK5 is involved with the development of opioid dependence. An animal study reported increased gene expression of *Grk5* in some brain regions in rats that were in a state of morphine withdrawal precipitated by naloxone (Fan et al., 2002).

Many heroin-dependent patients are also infected with HCV (Batki et al., 2010; Wang et al., 2011). In this study, the HCV infection rate was 94.6% in 344 MMT patients (Table 1), 91% in the 63 selected age- and gender-matched MMT patients, 63.6% in the 23 former heroin abusers, and 0% in normal controls (Table 2). It is known that HCV infection may influence the serum levels of AST and ALT (Crofts et al., 1993; Zechini et al., 2004; Aragon and Younossi, 2010). This could explain why the levels of AST and ALT were higher in MMT patients and former heroin abusers than the normal controls in the gene expression study (Table 2).

Limitations of this study include that all the identified genes did not pass the statistically significant threshold set for GWAS ($P < 3.2 \times 10^{-6}$). This indicated that the sample size in this study remains small for GWAS, which is common in pharmacogenomics studies (Motsinger-Reif et al., 2013). Gene expression was analyzed indirectly from EBV-transformed lymphoblastoid. Also, neuropsychological measures, for example, brain imaging, etc., are not available in this study. This precluded further explanations for the consequence of neurological changes after GRK5 expression alternations. Most subjects in this cohort were male, and 95% of these patients tested positive for HCV. In addition, the study was a cross-sectional design. However, participants in the study were recruited from a well-characterized, steady-state MMT cohort (Wang et al., 2013a). Medication compliance issues were well controlled by recruiting only subjects who had continuously undergone the MMT and shown regular attendance at the clinic. As the GRK5 gene has been reported to interact with multiple receptors and signal transduction pathways (Watari

et al., 2014), future studies should be conducted in search of pathways involving the methadone dose. Studies attempting to replicate the results and explore neuronal mechanisms of GRK5 for methadone treatment and opioid use are in need.

In summary, we identified that SNPs and haplotypes in GRK5 were associated with methadone dose in this study. When the levels of GRK5 gene expression were compared among 3 independent age- and gender-matched cohorts, MMT patients and medication-free former heroin abusers showed a higher level of gene expression than normal controls. These results suggested the involvement of GRK5 with the regulatory mechanisms of methadone dose and course of heroin dependence.

Supplementary Materials

Supplementary data are available at: *International Journal of Neuropsychopharmacology* online.

Funding

This work was supported by grants from the National Research Program for Genomic Medicine (NSC 100-3112-B-400-015); National Science Council (NSC 100-2314-B-400-002-MY3); Ministry of Science and Technology (MOST 106-2320-B-400-012) and the National Health Research Institutes, Taiwan (NP-106-PP-06, NP-107-PP-06, NP-106-SP-04 and NP-107-SP-04).

Acknowledgments

We acknowledge staff at the National Center for Genomic Medicine at Academia Sinica for their assistance in genotyping. We also thank staff in the Clinical Trial Information Management System (CTIMEs) at the National Health Research Institutes for data collection and management.

Statement of Interest

None.

References

- Aragon G, Younossi ZM (2010) When and how to evaluate mildly elevated liver enzymes in apparently healthy patients. *Cleve Clin J Med* 77:195–204.
- Arawaka S, et al (2006) The role of G-protein-coupled receptor kinase 5 in pathogenesis of sporadic Parkinson's disease. *J Neurosci* 26:9227–9238.
- Barratt DT, Collier JK, Hallinan R, Byrne A, White JM, Foster DJ, Somogyi AA (2012) ABCB1 haplotype and OPRM1 118A > G genotype interaction in methadone maintenance treatment pharmacogenetics. *Pharmacogenomics Pers Med* 5:53–62.
- Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21:263–265.
- Batki SL, Canfield KM, Smyth E, Ploutz-Snyder R, Levine RA (2010) Hepatitis C treatment eligibility and comorbid medical illness in methadone maintenance (MMT) and non-MMT patients: a case-control study. *J Addict Dis* 29:359–369.
- Chan YK, Sim DSM (2015) Steady-state principles. In: *Pharmacological Basis of Acute Care* (Chan KY, Ng PK, Sim MDS, eds), pp49–55. Cham, Switzerland: Springer International Publishing.
- Collier JK, Barratt DT, Dahlen K, Loennechen MH, Somogyi AA (2006) ABCB1 genetic variability and methadone dosage requirements in opioid-dependent individuals. *Clin Pharmacol Ther* 80:682–690.
- Crofts N, Hopper JL, Bowden DS, Breschkin AM, Milner R, Locarnini SA (1993) Hepatitis C virus infection among a cohort of victorian injecting drug users. *Med J Aust* 159:237–241.
- Devlin B, Roeder K (1999) Genomic control for association studies. *Biometrics* 55:997–1004.
- Fan X, Zhang J, Zhang X, Yue W, Ma L (2002) Acute and chronic morphine treatments and morphine withdrawal differentially regulate GRK2 and GRK5 gene expression in rat brain. *Neuropharmacology* 43:809–816.
- Fonseca F, de la Torre R, Díaz L, Pastor A, Cuyàs E, Pizarro N, Khymenets O, Farré M, Torrens M (2011) Contribution of cytochrome P450 and ABCB1 genetic variability on methadone pharmacokinetics, dose requirements, and response. *Plos One* 6:e19527.
- Glück L, Loktev A, Moulédous L, Mollereau C, Law PY, Schulz S (2014) Loss of morphine reward and dependence in mice lacking G protein-coupled receptor kinase 5. *Biol Psychiatry* 76:767–774.
- Hiltunen AJ, Eklund C (2002) Withdrawal from methadone maintenance treatment. Reasons for not trying to quit methadone. *Eur Addict Res* 8:38–44.
- Hung CC, Chiou MH, Huang BH, Hsieh YW, Hsieh TJ, Huang CL, Lane HY (2011) Impact of genetic polymorphisms in ABCB1, CYP2B6, OPRM1, ANKK1 and DRD2 genes on methadone therapy in han chinese patients. *Pharmacogenomics* 12:1525–1533.
- Jakobsdottir J, Conley YP, Weeks DE, Mah TS, Ferrell RE, Gorin MB (2005) Susceptibility genes for age-related maculopathy on chromosome 10q26. *Am J Hum Genet* 77:389–407.
- Levrano O, O'Hara K, Peles E, Li D, Barral S, Ray B, Borg L, Ott J, Adelson M, Kreek MJ (2008) ABCB1 (MDR1) genetic variants are associated with methadone doses required for effective treatment of heroin dependence. *Hum Mol Genet* 17:2219–2227.
- Levrano O, Peles E, Hamon S, Randesi M, Adelson M, Kreek MJ (2013a) CYP2B6 snps are associated with methadone dose required for effective treatment of opioid addiction. *Addict Biol* 18:709–716.
- Levrano O, Peles E, Randesi M, Shu X, Ott J, Shen PH, Adelson M, Kreek MJ (2013b) Association of genetic variation in pharmacodynamic factors with methadone dose required for effective treatment of opioid addiction. *Pharmacogenomics* 14:755–768.
- Li MX, Gui HS, Kwan JS, Sham PC (2011) GATES: a rapid and powerful gene-based association test using extended simes procedure. *Am J Hum Genet* 88:283–293.
- Li MX, Kwan JS, Sham PC (2012) HYST: a hybrid set-based test for genome-wide association studies, with application to protein-protein interaction-based association analysis. *Am J Hum Genet* 91:478–488.
- Mann A, Illing S, Miess E, Schulz S (2015) Different mechanisms of homologous and heterologous μ -opioid receptor phosphorylation. *Br J Pharmacol* 172:311–316.
- Marsden J, Farrell M, Bradbury C, Dale-Perera A, Eastwood B, Roxburgh M, Taylor S (2008) Development of the treatment outcomes profile. *Addiction* 103:1450–1460.
- Mattick RP, Breen C, Kimber J, Davoli M (2002) Methadone maintenance therapy vs no opioid replacement therapy for opioid dependence. *Cochrane Database Syst Rev* CD002209.
- Mattick RP, Breen C, Kimber J, Davoli M (2003) Methadone maintenance therapy vs no opioid replacement therapy for opioid dependence. *Cochrane Database Syst Rev* CD002209.

- Mattick RP, Breen C, Kimber J, Davoli M (2009) Methadone maintenance therapy vs no opioid replacement therapy for opioid dependence. *Cochrane Database Syst Rev* CD002209.
- Motsinger-Reif AA, Jorgenson E, Relling MV, Kroetz DL, Weinshilboum R, Cox NJ, Roden DM (2013) Genome-wide association studies in pharmacogenomics: successes and lessons. *Pharmacogenet Genomics* 23:383–394.
- Mouly S, Bloch V, Peoc'h K, Houze P, Labat L, Ksouda K, Simoneau G, Declèves X, Bergmann JF, Scherrmann JM, Laplanche JL, Lepine JP, Vorspan F (2015) Methadone dose in heroin-dependent patients: role of clinical factors, comedications, genetic polymorphisms and enzyme activity. *Br J Clin Pharmacol* 79:967–977.
- Murray H, McHugh RK, Behar E, Pratt E, Otto M (2008) Personality factors associated with methadone maintenance dose. *Am J Drug Alcohol Abuse* 34:634–641.
- Pitcher JA, Freedman NJ, Lefkowitz RJ (1998) G protein-coupled receptor kinases. *Annu Rev Biochem* 67:653–692.
- Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, Boehnke M, Abecasis GR, Willer CJ (2010) LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* 26:2336–2337.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81:559–575.
- Rodi C, Darnhofer-Patel B, Stanssens P, Zabeau M, van den Boom D (2002) A strategy for the rapid discovery of disease markers using the MassARRAY system. *BioTechniques* 32:S62–S69.
- Smith AH, Jensen KP, Li J, Nunez Y, Farrer LA, Hakonarson H, Cook-Sather SD, Kranzler HR, Gelernter J (2017) Genome-wide association study of therapeutic opioid dosing identifies a novel locus upstream of OPRM1. *Mol Psychiatry* 22:346–352.
- Storey J (2003) The positive false discovery rate: a bayesian interpretation and the q-value. *Ann Stat* 31:2013–2035.
- Sullivan SG, Wu Z, Cao X, Liu E, Detels R, National Methadone Maintenance Treatment Working Group (2014) Continued drug use during methadone treatment in china: a retrospective analysis of 19,026 service users. *J Subst Abuse Treat* 47:86–92.
- Tarantino P, De Marco EV, Annesi G, Rocca FE, Annesi F, Civitelli D, Provenzano G, Scornaienchi V, Greco V, Colica C, Nicoletti G, Quattrone A (2011) Lack of association between G-protein coupled receptor kinase 5 gene and Parkinson's disease. *Am J Med Genet B Neuropsychiatr Genet* 156B:104–107.
- Tompkins DA, Bigelow GE, Harrison JA, Johnson RE, Fudala PJ, Strain EC (2009) Concurrent validation of the clinical opiate withdrawal scale (COWS) and single-item indices against the clinical institute narcotic assessment (CINA) opioid withdrawal instrument. *Drug Alcohol Depend* 105:154–159.
- Tost J, Gut IG (2005) Genotyping single nucleotide polymorphisms by MALDI mass spectrometry in clinical applications. *Clin Biochem* 38:335–350.
- Tsai HJ, Wang SC, Liu SW, Ho IK, Chang YS, Tsai YT, Lin KM, Liu YL (2014) Assessment of CYP450 genetic variability effect on methadone dose and tolerance. *Pharmacogenomics* 15:977–986.
- Wang SC, et al. (2012) Genetic polymorphisms in the opioid receptor mu1 gene are associated with changes in libido and insomnia in methadone maintenance patients. *Eur Neuropsychopharmacol* 22:695–703.
- Wang SC, Tsou HH, Ho IK, Lin KM, Liu YL (2013a) Pharmacogenomics study in a Taiwan methadone maintenance cohort. *J Food Drug Anal* 21:S62–S68.
- Wang SC, Ho IK, Tsou HH, Liu SW, Hsiao CF, Chen CH, Tan HK, Lin L, Wu CS, Su LW, Huang CL, Yang YH, Liu ML, Lin KM, Liu SC, Wu HY, Kuo HW, Chen AC, Chang YS, Liu YL (2013b) Functional genetic polymorphisms in CYP2C19 gene in relation to cardiac side effects and treatment dose in a methadone maintenance cohort. *Omics* 17:519–526.
- Wang X, Zhang T, Ho WZ (2011) Opioids and HIV/HCV infection. *J Neuroimmune Pharmacol* 6:477–489.
- Watari K, Nakaya M, Kurose H (2014) Multiple functions of G protein-coupled receptor kinases. *J Mol Signal* 9:1.
- Wesson DR, Ling W (2003) The clinical opiate withdrawal scale (COWS). *J Psychoactive Drugs* 35:253–259.
- Yang HC, Chu SK, Huang CL, Kuo HW, Wang SC, Liu SW, Ho IK, Liu YL (2016) Genome-wide pharmacogenomic study on methadone maintenance treatment identifies SNP rs17180299 and multiple haplotypes on CYP2B6, SPON1, and GSG1L associated with plasma concentrations of methadone R- and S-enantiomers in heroin-dependent patients. *Plos Genet* 12:e1005910.
- Yuan HY, Chiou JJ, Tseng WH, Liu CH, Liu CK, Lin YJ, Wang HH, Yao A, Chen YT, Hsu CN (2006) FASTSNP: an always up-to-date and extendable service for SNP function analysis and prioritization. *Nucleic Acids Res* 34:W635–W641.
- Zechini B, Pasquazzi C, Aceti A (2004) Correlation of serum aminotransferases with HCV RNA levels and histological findings in patients with chronic hepatitis C: the role of serum aspartate transaminase in the evaluation of disease progression. *Eur J Gastroenterol Hepatol* 16:891–896.