

2019

Erratum: Best practices for evaluation and treatment of agitated children and adolescents (Beta) in the emergency department: Consensus statement of the American Association for Emergency psychiatry (Western Journal of Emergency Medicine (2019) 20:2 (409-419) DOI: 10.5811/westjem.2019.1.41344)

R. Gerson

N. Malas

V. Feuer

Zucker School of Medicine at Hofstra/Northwell

G. H. Silver

R. Prasad

See next page for additional authors

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>

 Part of the [Psychiatry Commons](#)

Recommended Citation

Gerson R, Malas N, Feuer V, Silver GH, Prasad R, Mroczkowski MM. Erratum: Best practices for evaluation and treatment of agitated children and adolescents (Beta) in the emergency department: Consensus statement of the American Association for Emergency psychiatry (Western Journal of Emergency Medicine (2019) 20:2 (409-419) DOI: 10.5811/westjem.2019.1.41344). . 2019 Jan 01; 20(3):Article 5782 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/5782>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

Authors

R. Gerson, N. Malas, V. Feuer, G. H. Silver, R. Prasad, and M. M. Mroczkowski

This Article Corrects: “Best Practices for Evaluation and Treatment of Agitated Children and Adolescents (BETA) in the Emergency Department: Consensus Statement of the American Association for Emergency Psychiatry”

Ruth Gerson, MD*
Nasuh Malas, MD, MPH†
Vera Feuer, MD‡
Gabrielle H. Silver, MD§
Raghuram Prasad, MD¶
Megan M. Mroczkowski, MD||

*Bellevue Hospital/New York University, Department of Psychiatry, New York, New York
†University of Michigan, Departments of Psychiatry and Pediatrics, Ann Arbor, Michigan
‡Northwell Health, Department of Psychiatry, New Hyde Park, New York
§Weill Cornell Medical College, Department of Psychiatry, New York, New York
¶Children’s Hospital of Philadelphia, Department of Psychiatry, Philadelphia, Pennsylvania
||Columbia University Medical Center, Department of Psychiatry, New York, New York

Electronically published February 19, 2019

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2019.4.43550

West J Emerg Med. 2019 March;20(2):409-419

Best Practices for Evaluation and Treatment of Agitated Children and Adolescents (BETA) in the Emergency Department: Consensus Statement of the American Association for Emergency Psychiatry
Gerson R, Malas N, Feuer V, Silver GH, Prasad R, Mroczkowski MM

Erratum in

West J Emerg Med. 2019 May;20(3):537. Pediatric BETA Consensus Guideline Working Group members should not be included in full author list. The author list has now been corrected on this erratum.

Abstract

Introduction: Agitation in children and adolescents in the emergency department (ED) can be dangerous and distressing for patients, family and staff. We present consensus guidelines for management of agitation among pediatric patients in the ED, including non-pharmacologic methods and the use of immediate and as-needed medications.

Methods: Using the Delphi method of consensus, a workgroup comprised of 17 experts in emergency child and adolescent psychiatry and psychopharmacology from the the American Association for Emergency Psychiatry and the American Academy of Child and Adolescent Psychiatry Emergency Child Psychiatry Committee sought to create consensus guidelines for the management of acute agitation in children and adolescents in the ED.

Results: Consensus found that there should be a multimodal approach to managing agitation in the ED, and that etiology of agitation should drive choice of treatment. We describe general and specific recommendations for medication use.

Conclusion: These guidelines describing child and adolescent psychiatry expert consensus for the management of agitation in the ED may be of use to pediatricians and emergency physicians who are without immediate access to psychiatry consultation.

PMCID: PMC6404720 [PubMed - indexed for MEDLINE]