

2020

Impact of the COVID-19 pandemic on endoscopy practice: results of a cross-sectional survey from the New York metropolitan area

S. H. Mahadev

O. S. Aroniadis

L. Barraza

E. Agarunov

A. J. Goodman

See next page for additional authors

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/articles>

Recommended Citation

Mahadev SH, Aroniadis OS, Barraza L, Agarunov E, Goodman AJ, Benias PC, Buscaglia JM, Gross SA, Kasmin FE, Cohen JJ, Carr-Locke DL, Greenwald DA, Mendelsohn RB, Sethi A, Gonda TA. Impact of the COVID-19 pandemic on endoscopy practice: results of a cross-sectional survey from the New York metropolitan area. . 2020 Jan 01; ():Article 6212 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/articles/6212>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.

Authors

S. H. Mahadev, O. S. Aroniadis, L. Barraza, E. Agarunov, A. J. Goodman, P. C. Benias, J. M. Buscaglia, S. A. Gross, F. E. Kasmin, J. J. Cohen, D. L. Carr-Locke, D. A. Greenwald, R. B. Mendelsohn, A. Sethi, and T. A. Gonda



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Impact of the COVID-19 pandemic on endoscopy practice: results of a cross-sectional survey from the New York metropolitan area



To the Editor:

We read with great interest the recent articles by Repici et al¹ and Soetikno et al² outlining the approaches taken in Italy and Hong Kong to adapting endoscopy practice to meet the new challenges of the COVID-19 pandemic. Because the virus has now moved to our shores, we sought to describe our experience of the impact of COVID-19 on the practice of endoscopy in the New York metropolitan area, which currently has the highest case burden in the world.

To provide a snapshot of endoscopy activities from the beginning to the peak of the pandemic in the New York metropolitan area, we distributed an electronic survey in

collaboration with, and with the support of, the New York Society of Gastroenterological Endoscopy (NYSGE). Surveys were distributed to physicians on March 30, 2020, and were returned over the following 7 days. The survey collected information regarding demographics, case volume, personal protective equipment (PPE) use, and endoscopist exposure to COVID-19. All responses were collected and analyzed anonymously.

A total of 69 endoscopists responded to the survey. Eight (12%) were fellows in training; 21 (30%) were aged over 60. The majority (59%) practiced in an academic hospital endoscopy unit. Forty-one percent performed advanced endoscopy. Most endoscopists spent 2 to 4 days a week in the endoscopy suite before the pandemic.

Since the onset of the COVID-19, the case volume was markedly reduced, with 71% of gastroenterologists reporting that they no longer regularly performed endoscopy at all, and 62% reporting no cases over the preceding 7




Figure 1. Average weekly endoscopy unit volume and universal personal protective equipment (PPE) use before (A) and since (B) the onset of the COVID-19 pandemic.

days. Widespread restrictions on cases to urgent (67%) or no outpatients at all (31%) were reported to have been instituted an average of 2 weeks before the survey. Weekly endoscopy volumes declined by 57% to 96% for all procedure types, with EGD and ERCP being the most frequently performed during the pandemic (Fig. 1A). Eight endoscopists (13%) reported having performed endoscopy on confirmed COVID-19+ patients. The majority (53%) of physicians reported they had been redeployed from usual duties to cover COVID-19+ hospital medicine services.

Universal N95 respiratory use for all cases since the onset of COVID-19 was reported by only 65%. Many endoscopists reported that the use of N95 masks was restricted to known or suspected COVID-19 cases because of limited availability (17%) or were not available at all (9%). Our data also show that overall PPE use changed significantly in comparison with pre-COVID practices (Fig. 1B). Testing patients for COVID-19 was not routine before endoscopy for most respondents (54%) at the time of this survey.

In conclusion, COVID-19 has had a drastic impact on the practice of endoscopy and procedure volumes in the New York metropolitan area. These numbers may provide an early estimate of the impact of this pandemic on GI practices. The initial experience of expansion of universal PPE and limited availability of PPE are notable, especially in light of joint gastroenterology society guidelines released on April 1 that recommend universal N95 mask use for all endoscopy team members.³ Although these data reflect the first weeks of a rapidly evolving pandemic in the United States, they may inform preparedness efforts in regions that anticipate, or are at an earlier phase of, the pandemic.

DISCLOSURE

All authors disclosed no financial relationships.

SriHari Mahadev, MD, MS

*NYP-Weill Cornell Medicine
New York*

Olga S. Aroniadis, MD

*Stony Brook University Hospital
Stony Brook*

Luis Barraza, MD

Emil Agarunov, MD

*NYP-Columbia University Irving Medical Center
New York*

Adam J. Goodman, MD

*NYU Langone Health
New York*

Petros C. Benias, MD

*Northwell Health
Manhasset*

Jonathan M. Buscaglia, MD

*Stony Brook University Hospital
Stony Brook*

Seth A. Gross, MD

*NYU Langone Health
New York*

Franklin E. Kasmin, MD

*Mount Sinai Hospital
New York*

Jonathan J. Cohen, MD

*NYU Langone Health
New York*

David L. Carr-Locke, MD

*NYP-Weill Cornell Medicine
New York*

David A. Greenwald, MD

*Mount Sinai Hospital
New York*

Robin B. Mendelsohn, MD

*Memorial Sloan Kettering Hospital
New York*

Amrita Sethi, MD

Tamas A. Gonda, MD, on behalf of NYSGE

research committee

*NYP-Columbia University Irving Medical Center
New York, New York, USA*

REFERENCES

1. Repici A, Maselli R, Colombo M, et al. Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc* 2020;92:192-7.
2. Soetikno R, Teoh AY, Kaltenbach T, et al. Considerations in performing endoscopy during the COVID-19 pandemic. *Gastrointest Endosc* 2020;92:176-83.
3. American Society for Gastrointestinal Endoscopy, et al. Joint gastroenterology society message: COVID-19 use of personal protective equipment in GI endoscopy. (Joint message.) *Gastrointest Endosc*. Epub 2020 April 2.

<http://dx.doi.org/10.1016/j.gie.2020.04.047>

COVID-19 testing before every endoscopy: Is India ready for prime time?



To the Editor:

The first case of SARS-CoV-2 infection in India was reported on January 30, 2020, from Kerala.¹ The disease since then has increased manifold to reach figures of over 45,000 infections across the country, making a significant impact on healthcare with drastic changes in clinical practice. Multiple society guidelines have been published since the outbreak of the virus, with a major focus on screening and precautions for patients undergoing endoscopy. Continuing hospital services in a