

2020

Management of upper GI bleeding in patients with COVID-19 pneumonia

K Cavaliere
Northwell Health

C Levine
Zucker School of Medicine at Hofstra/Northwell, clevine5@northwell.edu

P Wander
Northwell Health

DV Sejpal
Zucker School of Medicine at Hofstra/Northwell, dsejpal@northwell.edu

AJ Trindade
Zucker School of Medicine at Hofstra/Northwell, atrindade@northwell.edu

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/articles>

 Part of the [Gastroenterology Commons](#)

Recommended Citation

Cavaliere K, Levine C, Wander P, Sejpal D, Trindade A. Management of upper GI bleeding in patients with COVID-19 pneumonia. . 2020 Jan 01; 92(2):Article 6390 [p.]. Available from: <https://academicworks.medicine.hofstra.edu/articles/6390>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact academicworks@hofstra.edu.



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

*Hubei Provincial Clinical Research Center
for Digestive Disease Minimally Invasive Incision
Renmin Hospital of Wuhan University
Wuhan, China*

Xuefen Wang, MS
*Nursing Department
Renmin Hospital of Wuhan University
Wuhan, China*

Zhongyin Zhou, MD
Honggang Yu, MD, PhD
*Department of Gastroenterology
Key Laboratory of Hubei Province for
Digestive System Disease
Hubei Provincial Clinical Research Center
for Digestive Disease Minimally Invasive Incision
Renmin Hospital of Wuhan University
Wuhan, China*

Drs An, Huang, and Wan contributed equally and are joint first authors.

Supported in part by grants from the National Natural Science Foundation of China (Grant No. 81672387 to Yu Honggang) and the National Natural Science Foundation of China (Grant No. 81302131 to Ping An).

REFERENCES

1. Repici A, Maselli R, Colombo M, et al. Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc.* Epub 2020 Mar 14.
2. Soetikno R, Teoh AY, Kaltenbach T, et al. Considerations in performing endoscopy during the COVID-19 pandemic. *Gastrointest Endosc.* Epub 2020 Mar 27.
3. The National Health and Health Commission of China. Available at: http://en.nhc.gov.cn/2020-02/02/c_76098.htm. Accessed February 2, 2020.
4. Beilenhoff U, Biering H, Blum R, et al. Reprocessing of flexible endoscopes and endoscopic accessories used in gastrointestinal endoscopy: position statement of the European Society of Gastrointestinal Endoscopy (ESGE) and European Society of Gastroenterology Nurses and Associates (ESGENA) - Update 2018. *Endoscopy* 2018;50:1205-34.
5. European Centre for Disease Prevention and Control. Personal protective equipment (PPE) needs in healthcare settings for the care of patients with suspected or confirmed novel coronavirus (2019-nCoV) 2020. Stockholm, Sweden: European Centre for Disease Prevention and Control; 2020.

<https://doi.org/10.1016/j.gie.2020.04.022>

Management of upper GI bleeding in patients with COVID-19 pneumonia



To the Editor:

Coronavirus disease 2019 (COVID-19) has become a worldwide pandemic. The typical presentation is a respiratory illness with fever, cough, and shortness of breath. GI symptoms are being increasingly recognized and include abdominal pain, vomiting, diarrhea, and nausea.¹ We present a case series of 6 patients who were admitted to our hospital with COVID-19-associated pneumonia (fever, shortness of breath requiring oxygen, positive COVID-19 polymerase chain reaction test result, and infil-

trates showing on chest radiograph), and upper GI bleeding. The patient and clinical characteristics are shown in Table 1. The GI manifestations were hematemesis or melena.

Guidelines advise that patients who present with acute upper GI bleeding undergo endoscopy within 24 hours of presentation.² Endoscopy can not only provide therapy but also allow for risk stratification for re-bleeding that can dictate management. However, the discussion for endoscopy in patients with COVID-19 pneumonia brings about unique management decisions. Although endoscopy can provide therapy if a discrete visible vessel is seen, the risk of the procedure may outweigh the benefit in patients with COVID-19 pneumonia.

First, 5 of the 6 patients in this series were receiving supplemental oxygen, and 1 patient had an endotracheal tube. Performing upper endoscopy would have likely required general anesthesia with an endotracheal tube in the 5 patients, given their oxygen requirements, the indication for the procedure (hematemesis), or both. Extubation after the procedure becomes challenging in the setting of pneumonia. In addition, a recent study from China demonstrated an increased mortality rate once a patient with COVID-19 pneumonia is intubated.³ Although the data for this concerned emergent intubation for respiratory failure (not an elective procedure), the data are compelling. Second, there is a real concern for transmission of the virus to the anesthesiologist, staff, and endoscopist, given the aerosolization of respiratory droplets during endoscopy.⁴

Given that the risks of endoscopy might outweigh the benefits, we decided to treat these patients conservatively with a proton pump inhibitor drip, blood transfusion as needed, and frequent monitoring of vital signs, GI symptoms, and hemoglobin value. Endoscopy was reserved if the patient did not respond to conservative management by 24 hours (lack of hemodynamic stability and if the hemoglobin was not stable). Delaying the endoscopy for 24 hours has recently been shown to not affect 30-day mortality in comparison with earlier endoscopy.⁵ All 6 of our patients responded to conservative management. Cessation of clinical symptoms of acute upper GI bleeding was seen in all of our patients in combination with stabilization of hemoglobin. None of the patients required upper endoscopy during their clinical course.

The exact cause of GI bleeding in this cohort is unknown because endoscopy was not performed. The most likely cause is ulcer related. Another cause being recognized is COVID-related coagulopathy.⁶ Given that the patients responded to conservative management, the former is more likely.

In conclusion, the treatment of patients admitted with COVID-19 pneumonia who experience upper GI bleeding is challenging. It can possibly be managed

TABLE 1. Case series of 6 patients with COVID-19 pneumonia and upper GI bleeding

Patient	Age, y	Gender	Presenting GI symptom	O ₂ saturation	Hb g/dL, HCT %, Plts K/uL	Transfusion	CXR finding	GBS	D-dimer, ng/mL*	Ferritin, ng/mL†	LDH, U/L‡	GI outcome
1	77	Male	Hematemesis	94% RA	9.6/30.5/154	No	BI	12	372	860	399	GI bleed resolved
2	65	Male	Melena	86% RA	6.2/17.7/151	Yes 2 U PRBC	BO	18	3042	7987	1142	GI bleed resolved
3	46	Male	Melena	90% RA	6.8/21.2/226	Yes 2 U PRBC	BI	11	285	3970	878	GI bleed resolved
4	70	Female	Melena	92% RA	10.2/31.2/260	No	BO	11	38674	1040	1188	GI bleed resolved
5	67	Female	Hematemesis	88% RA	6.1/20.7/209	Yes 2 U PRBC	BI RUL opacity	15	-	802	563	GI bleed resolved
6	82	Female	Melena	94% 5LO ₂	8.6/29.2/302	Yes 2 U PRBC	BI LLL opacity	14	1198	705	959	GI bleed resolved

RA, Room air; CXR, chest x-ray; GBS, Glasgow-Blatchford score; BI, bilateral interstitial infiltrates; BO, bilateral opacities; RUL, right upper lobe; LLL, left lower lobe; Hb, hemoglobin; HCT, hematocrit; LDH, lactate dehydrogenase; PRBC, packed red blood cells; Plts, platelets; 5LO₂, five liters oxygen.

*D-dimer: upper limit of normal <229 ng/mL.

†Ferritin: upper limit of normal 400 ng/mL.

‡LDH: upper limit of normal 242 U/L.

conservatively without endoscopy because all of our patients responded by 24 hours. Lack of response in 24 hours may indicate a need for endoscopy with personal protective equipment.

DISCLOSURE

Dr Sejpal is a consultant for Boston Scientific, Olympus America, and Gyrus ACMI, Inc. Dr Trindade is a consultant for Olympus America and Pentax Medical, the recipient of research support from Ninepoint Medical, and an attendee at educational events by Boston Scientific. The other authors disclosed no financial relationships.

Kimberly Cavaliere, MD
Calley Levine, MD
Praneet Wander, MBBS
Divyesh V. Sejpal, MD
Arvind J. Trindade, MD

*Division of Gastroenterology
 Long Island Jewish Medical Center
 Zucker School of Medicine at Hofstra/Northwell
 Northwell Health System
 New Hyde Park, New York, USA*

REFERENCES

- Zhou Z, Zhao N, Shu Y, et al. Effect of gastrointestinal symptoms on patients infected with COVID-19. *Gastroenterology*. Epub 2020 Mar 18.
- Laine L, Jensen DM. Management of patients with ulcer bleeding. *Am J Gastroenterol* 2012;107:345-60.
- Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. Epub 2020 Feb 24.

- Marchese M, Capannolo A, Lombardi L, et al. Use of a modified ventilation mask to avoid aerosolizing spread of droplets for short endoscopic procedures during coronavirus COVID-19 outbreak. *Gastrointest Endosc*. Epub 2020 Apr 2.
- Lau JYW, Yu Y, Tang RSY, et al. Timing of endoscopy for acute upper gastrointestinal bleeding. *N Engl J Med* 2020;382:1299-308.
- Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost*. Epub 2020 Mar 25.

<https://doi.org/10.1016/j.gie.2020.04.028>

Dose response for argon plasma coagulation in the treatment of weight regain after Roux-en-Y gastric bypass: Is a high dose really necessary?



To the Editor:

We read with interest the recent article by Jirapinyo et al¹ regarding the dose response for argon plasma coagulation (APC) in the treatment of weight regain after Roux-en-Y gastric bypass. Indeed, it is a practical article assessing the outcomes of different electro-surgery settings for ablating the dilated gastrojejunostomy. The authors report more significant weight loss and a similar safety profile with the high-dose APC setting. However, some concerning methodologic aspects drew our attention, and we would like to share them.

First, the authors report in their description of the demographics that 31% of patients had undergone transoral outlet reduction before the APC procedure. From that, we take it that those patients probably received both ablation and suturing at the anastomosis. However, there is no mention of the distribution of the previous transoral