

2020

## Intensive care unit admissions for pregnant and nonpregnant women with coronavirus disease 2019

M J. Blitz

*Zucker School of Medicine at Hofstra/Northwell*

A Grünebaum

*Zucker School of Medicine at Hofstra/Northwell*

A Tekbali

*Northwell Health*

E Bornstein

*Zucker School of Medicine at Hofstra/Northwell*

B Rochelson

*Zucker School of Medicine at Hofstra/Northwell*

*See next page for additional authors*

Follow this and additional works at: <https://academicworks.medicine.hofstra.edu/publications>



Part of the [Obstetrics and Gynecology Commons](#)

---

### Recommended Citation

Blitz MJ, Grünebaum A, Tekbali A, Bornstein E, Rochelson B, Nimaroff M, Chervenak F. Intensive care unit admissions for pregnant and nonpregnant women with coronavirus disease 2019. . 2020 Jan 01; 223(2):Article 6465 [ p.]. Available from: <https://academicworks.medicine.hofstra.edu/publications/6465>. Free full text article.

This Article is brought to you for free and open access by Donald and Barbara Zucker School of Medicine Academic Works. It has been accepted for inclusion in Journal Articles by an authorized administrator of Donald and Barbara Zucker School of Medicine Academic Works. For more information, please contact [academicworks@hofstra.edu](mailto:academicworks@hofstra.edu).

---

**Authors**

M J. Blitz, A Grünebaum, A Tekbali, E Bornstein, B Rochelson, M Nimaroff, and FA Chervenak



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obstet Gynecol MFM* 2020. [Epub ahead of print].

2. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020. [Epub ahead of print].

3. Dashraath P, Wong JLJ, Lim MXK, et al. Coronavirus disease 2019 (COVID-19) pandemic and pregnancy. *Am J Obstet Gynecol* 2020. [Epub ahead of print].

4. Mullins E, Evans D, Viner RM, O'Brien P, Morris E. Coronavirus in pregnancy and delivery: rapid review. *Ultrasound Obstet Gynecol* 2020. [Epub ahead of print].

5. Zaigham M, Andersson O. Maternal and perinatal outcomes with COVID-19: a systematic review of 108 pregnancies. *Acta Obstet Gynecol Scand* 2020. [Epub ahead of print].

6. NIH. COVID-19 treatment guidelines. Available at: <https://covid19.treatmentguidelines.nih.gov/introduction/>. Accessed April 23, 2020.

© 2020 Elsevier Inc. All rights reserved. <https://doi.org/10.1016/j.ajog.2020.04.029>

## Intensive care unit admissions for pregnant and nonpregnant women with coronavirus disease 2019



**OBJECTIVE:** Early reports indicate that pregnant women are not at an increased risk for coronavirus disease 2019 (COVID-19) or for a worse disease course if infection occurs.<sup>1-3</sup> This study aimed to review our experiences with intensive care unit (ICU) admissions of women of reproductive age with COVID-19, and to determine whether pregnant women are more likely to be admitted to the ICU than nonpregnant women.

**STUDY DESIGN:** We evaluated data from a large hospital system in New York State between March 2, 2020, and April 9, 2020. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) testing was performed on acutely symptomatic patients presenting with characteristic respiratory signs and symptoms.<sup>4</sup> Nasopharyngeal specimens were obtained, and microbiologic diagnosis was made on the basis of a positive result on SARS-CoV-2 real-time reverse transcription polymerase chain reaction (RT-PCR) assay. We included only patients of reproductive age (15–49 years) who were admitted to 1 of 7 hospitals in our system and who received a diagnosis of COVID-19 by RT-PCR on

admission, during the hospital stay, or during the postpartum period. Data analyzed in this study included age (5 age groups between 15 and 49 years), pregnancy status, and admission to an ICU. We excluded patients with incomplete data. The incidence of ICU admission was compared between pregnant and nonpregnant women with COVID-19 in each age group. Patients were admitted to the ICU at the discretion of the consulted critical care attending physician. Other clinical characteristics, including medical comorbidities, were not evaluated and not necessarily the same. The institutional review board determined that this study did not meet the definition of human subjects research and was exempt from formal review.

**RESULTS:** Among all patients between the ages of 15–49 years admitted at 7 hospitals within our health system between March 4, 2020, and April 9, 2020, there were 1168 symptomatic patients who received a diagnosis of COVID-19. Of these, 754 (64.6%) were male, 332 (28.4%) were nonpregnant females, and 82 (7.0%) were pregnant females. During this time period, 2971 pregnant patients were admitted, primarily

**TABLE**

**ICU admissions by age group in pregnant and nonpregnant women with COVID-19**

Age group, y	Pregnant women (n = 82)	Nonpregnant women (n = 332)	P value
<25	1/11 (9.1)	3/7 (42.9)	.09
25–29	0/17 (0)	5/40 (12.5)	.16
30–34	2/33 (6.1)	5/44 (11.4)	.46
35–39	3/15 (20.0)	9/55 (16.4)	.73
40–49	2/6 (33.3)	28/190 (14.7)	.28
Total	8/82 (9.8)	50/332 (15.1)	.22

Values are presented as n/N (%).

COVID-19, coronavirus disease 2019; ICU, intensive care unit.

Blitz. ICU admissions for pregnant and nonpregnant women with COVID-19. *Am J Obstet Gynecol* 2020.

for delivery. In some cases, symptomatic patients with a diagnosis of COVID-19 (2.8%) were admitted for obstetrical indications and only had mild respiratory disease. In all, 50 nonpregnant females (15.1%, 50 of 332) and 8 pregnant females (9.8%, 8 of 82) were admitted to the ICU for worsening respiratory status, with no statistically significant difference ( $P=.22$ ). ICU admissions by age group are presented in the [Table](#).

**CONCLUSION:** Among hospitalized women who received a diagnosis of COVID-19, pregnant women were not at an increased risk for ICU admission compared with nonpregnant women. This finding is consistent with the overall lower hospital admission rate of pregnant women with COVID-19 that we previously reported.<sup>5</sup> Pregnant women are considered to be at a greater risk of severe morbidity and mortality from other respiratory infections such as influenza.<sup>6</sup> Admission to the ICU signifies a more severe course of disease. Therefore, our findings are reassuring and indicate that pregnant women with COVID-19 may not experience more severe disease progression than nonpregnant women. ■

Matthew J. Blitz, MD, MBA  
Donald and Barbara Zucker School of Medicine at Hofstra/Northwell  
376 East Main Street  
Suite 202  
Bay Shore, NY 11706  
[mblitz@northwell.edu](mailto:mblitz@northwell.edu)

Amos Grünebaum, MD  
Asma Tekbali, MPH

Eran Bornstein, MD  
Burton Rochelson, MD  
Michael Nimaroff, MD  
Frank A. Chervenak, MD  
Donald and Barbara Zucker School of Medicine at Hofstra/Northwell  
Manhasset, NY

The authors report no conflict of interest.

This communication has been published in the middle of the COVID-19 pandemic and is available via expedited publication to assist patients and healthcare providers.

#### REFERENCES

1. Rasmussen SA, Smulian JC, Lednicky JA, Wen TS, Jamieson DJ. Coronavirus disease 2019 (COVID-19) and pregnancy: what obstetricians need to know. *Am J Obstet Gynecol* 2020;222:415–26.
2. Della Gatta AN, Rizzo R, Pilu G, Simonazzi G. COVID19 during pregnancy: a systematic review of reported cases. *Am J Obstet Gynecol* 2020. [Epub ahead of print].
3. Yan J, Guo J, Fan C, et al. Coronavirus disease 2019 (COVID-19) in pregnant women: a report based on 116 cases. *Am J Obstet Gynecol* 2020. [Epub ahead of print].
4. Centers for Disease Control and Prevention. Symptoms of coronavirus. 2020. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>. Accessed April 27, 2020.
5. Tekbali A, Grünebaum A, Saraya A, McCullough L, Bornstein E, Chervenak FA. Pregnant versus non-pregnant SARS-CoV-2 and COVID-19 hospital admissions: the first 4 weeks in New York. *Am J Obstet Gynecol* 2020. [Epub ahead of print].
6. Sappenfield E, Jamieson DJ, Kourtis AP. Pregnancy and susceptibility to infectious diseases. *Infect Dis Obstet Gynecol* 2013;2013:752852.

© 2020 Elsevier Inc. All rights reserved. <https://doi.org/10.1016/j.ajog.2020.05.004>

## Universal severe acute respiratory syndrome coronavirus 2 testing of pregnant women admitted for delivery in 2 Italian regions



**BACKGROUND:** Since the early days of the coronavirus disease 2019 (COVID-19) pandemic, substantial undocumented infection has been thought to contribute to the dissemination of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2),<sup>1</sup> with estimated percentages of undocumented infections of 72%–90%.<sup>1–3</sup>

Italy has been the first western country to be massively hit by the pandemic. On March 11, the Italian government ordered the country lockdown, which is still in place as of this writing. This led to a flattening and eventually a reduction of the pandemic curve.

Starting by the end of March, several hospitals have begun universal SARS-CoV-2 screening in all admitted patients. Women admitted for delivery represent a peculiar population and a unique source of information, because

they come to hospitals independent of illness and of their own decision. They can therefore provide useful estimates of the circulation of the virus in the general population, despite a possible different social behavior, especially near delivery.

**OBJECTIVE:** This study aimed to estimate the “true” SARS-CoV-2 infection rate among women admitted for delivery and estimate the burden of undocumented infections in this population.

**METHODS:** We studied 2 neighboring Italian regions, North of Tuscany and Liguria, both considered at medium risk of infection compared with the Northern regions. All 6 hospitals of Azienda USL “Toscana Nord Ovest” (ATNO, Tuscany) and